

$\eta_c(2S)$ 

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

 $\eta_c(2S)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3637.5 ± 1.1 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
3635.1 ± 3.7 ± 2.9	106	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
3633.6 ± 1.7 ± 0.6	106	<sup>1</sup> AAIJ	17ADLHCB	$pp \rightarrow B^+X \rightarrow p\bar{p}K^+X$
3636.4 ± 4.1 ± 0.7	365	<sup>2</sup> AAIJ	17BBLHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3637.0 ± 5.7 ± 3.4	178	<sup>3,4</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
3635.1 ± 5.8 ± 2.1	47	<sup>3,5</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
3646.9 ± 1.6 ± 3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127 ± 18	<sup>6</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, KK\pi^0$
3638.5 ± 1.5 ± 0.8	624	<sup>3</sup> DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	<sup>3</sup> DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
3636.1 <sup>+3.9+0.7</sup> <sub>-4.2-2.0</sub>	128	<sup>7</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	<sup>8</sup> ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3645.0 ± 5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05C BABR	$e^+e^- \rightarrow J/\psi c\bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3639 ± 7	98 ± 52	<sup>9</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_c \bar{c}$
3630.8 ± 3.4 ± 1.0	112 ± 24	<sup>10</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ± 6 ± 8	39 ± 11	<sup>11</sup> CHOI	02 BELL	$B \rightarrow K K_S K^-\pi^+$
3594 ± 5		<sup>12</sup> EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

<sup>1</sup>AAIJ 17AD report  $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$  MeV. We use the current value  $m_{\psi(2S)} = 3686.097 \pm 0.025$  MeV to obtain the quoted mass.

<sup>2</sup>From a fit of the  $\phi\phi$  invariant mass with the width of  $\eta_c(2S)$  fixed to the PDG 16 value.

<sup>3</sup>Ignoring possible interference with continuum.

<sup>4</sup>With a width fixed to 11.3 MeV.

<sup>5</sup>With a width fixed to 11.3 MeV. Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$  decays.

<sup>6</sup>From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+K^-\pi^0$  decay modes.

<sup>7</sup>Accounts for interference with non-resonant continuum.

<sup>8</sup>From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>9</sup>From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>10</sup>Superseded by DEL-AMO-SANCHEZ 11M.

<sup>11</sup>Superseded by VINOKUROVA 11.

<sup>12</sup>Assuming mass of  $\psi(2S) = 3686$  MeV.

$\eta_c(2S)$  WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.3<sup>+3.2</sup><sub>-2.9</sub></b>					<b>OUR AVERAGE</b>
9.9 <sup>±</sup> 4.8 <sup>±</sup> 2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 <sup>±</sup> 6.4 <sup>±</sup> 4.8		127 ± 18	<sup>13</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
13.4 <sup>±</sup> 4.6 <sup>±</sup> 3.2		624	<sup>14</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 <sup>+8.4+2.6</sup> <sub>-5.1-0.9</sub>		128	<sup>15</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 <sup>±</sup> 12.4 <sup>±</sup> 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	98 ± 52	<sup>16</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
17.0 <sup>±</sup> 8.3 <sup>±</sup> 2.5		112 ± 24	<sup>17</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
<55	90	39 ± 11	<sup>18</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
<8.0	95		<sup>19</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>13</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>14</sup> Ignoring possible interference with continuum.

<sup>15</sup> Accounts for interference with non-resonant continuum.

<sup>16</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>17</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>18</sup> For a mass value of  $3654 \pm 6$  MeV. Superseded by VINOKUROVA 11.

<sup>19</sup> For a mass value of  $3594 \pm 5$  MeV

 $\eta_c(2S)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	( 1.9 <sup>±</sup> 1.2 ) %	
$\Gamma_3$ $K\bar{K}\eta$	( 5 ± 4 ) × 10 <sup>-3</sup>	
$\Gamma_4$ $2\pi^+ 2\pi^-$	not seen	
$\Gamma_5$ $\rho^0 \rho^0$	not seen	
$\Gamma_6$ $3\pi^+ 3\pi^-$	not seen	
$\Gamma_7$ $K^+ K^- \pi^+ \pi^-$	not seen	
$\Gamma_8$ $K^{*0} \bar{K}^{*0}$	not seen	
$\Gamma_9$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.4 <sup>±</sup> 1.0 ) %	
$\Gamma_{10}$ $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
$\Gamma_{11}$ $K_S^0 K^- 2\pi^+ \pi^- + c.c.$	seen	
$\Gamma_{12}$ $2K^+ 2K^-$	not seen	
$\Gamma_{13}$ $\phi\phi$	not seen	

$\Gamma_{14}$	$p\bar{p}$	seen		
$\Gamma_{15}$	$p\bar{p}\pi^+\pi^-$	seen		
$\Gamma_{16}$	$\gamma\gamma$	$(1.9\pm 1.3)\times 10^{-4}$		
$\Gamma_{17}$	$\gamma J/\psi(1S)$	$< 1.4$	%	90%
$\Gamma_{18}$	$\pi^+\pi^-\eta$	not seen		
$\Gamma_{19}$	$\pi^+\pi^-\eta'$	not seen		
$\Gamma_{20}$	$\pi^+\pi^-\eta_c(1S)$	$< 25$	%	90%

### $\eta_c(2S)$ PARTIAL WIDTHS

#### $\Gamma(\gamma\gamma)$ $\Gamma_{16}$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.44\pm 0.14$	106	<sup>20</sup> XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$1.3\pm 0.6$		<sup>21</sup> ASNER	04	CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>20</sup> Assuming that the branching fraction into  $\eta'\pi^+\pi^-$  is the same as for  $\eta_c(1S)$ .

<sup>21</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV.

#### $\Gamma(\gamma\gamma) \times \Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_{19}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$5.6_{-1.1}^{+1.2} \pm 1.1$	106	XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
-----------------------------	-----	----	----	---

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

#### $\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$< 6.5$	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$
---------	----	--------	----	--

#### $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{16}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$41 \pm 4 \pm 6$	624	<sup>22</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
------------------	-----	-------------------------------	------	--

<sup>22</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

#### $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$< 5.0$	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$
---------	----	--------	----	---

#### $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_9\Gamma_{16}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$30 \pm 6 \pm 5$	1201	<sup>23</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
------------------	------	-------------------------------	------	--

<sup>23</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{12}\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{20}\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<133	90	LEES	12AE	BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$

VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90 <sup>24,25,26</sup>	AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90 <sup>24,25,27</sup>	AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	25,27	AMBROGIANI	01 E835 $\bar{p}p \rightarrow \gamma\gamma$

<sup>24</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>25</sup> For a total width  $\Gamma=5$  MeV.

<sup>26</sup> For the resonance mass region 3589–3599 MeV/ $c^2$ .

<sup>27</sup> For the resonance mass region 3575–3660 MeV/ $c^2$ .

$\eta_c(2S)$  BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABREU	98O	DLPH $e^+e^- \rightarrow e^+e^- + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>28</sup> EDWARDS	82C	CBAL $e^+e^- \rightarrow \gamma X$
------	-----------------------	-----	------------------------------------

<sup>28</sup> For a mass value of  $3594 \pm 5$  MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.4 \pm 1.1$	$59 \pm 12$	<sup>29</sup> AUBERT	08AB	BABR $B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	$127 \pm 18$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
------	--------------	---------	-----	--

seen	$39 \pm 11$	<sup>30</sup> CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$
------	-------------	--------------------	----	--------------------------------------

<sup>29</sup> Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .

<sup>30</sup> For a mass value of  $3654 \pm 6$  MeV

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$   $\Gamma_3/\Gamma_2$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$27.3 \pm 7.0 \pm 9.0$	225	<sup>31</sup> LEES	14E	BABR $\gamma\gamma \rightarrow K^+K^-\gamma\gamma$

<sup>31</sup> LEES 14E reports  $B(\eta_c(2S) \rightarrow K^+K^-\eta)/B(\eta_c(2S) \rightarrow K^+K^-\pi^0) = 0.82 \pm 0.21 \pm 0.27$ , which we divide by 3 to account for isospin symmetry.

$\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$   $\Gamma_9/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.73 \pm 0.17 \pm 0.17$	1201	<sup>32</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

<sup>32</sup>We have multiplied the value of  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K_S^0K^\pm\pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+K^-\pi^+\pi^-$

$\Gamma(K_S^0K^-2\pi^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	$57 \pm 17$	ABLIKIM 13K	BES3	$\psi(2S) \rightarrow \gamma K_S^0K^\pm\pi^\mp\pi^+\pi^-$

$\Gamma(2K^+2K^-)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+K^-K^+K^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	106	<sup>33</sup> AAIJ 17AD	LHCB	$pp \rightarrow B^+X \rightarrow p\bar{p}K^+X$

<sup>33</sup>AAIJ 17AD report a 6.4 standard deviation signal, with  $B(B^+ \rightarrow \eta_c(2S)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$ .

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	110	<sup>34</sup> CHILIKIN 19	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>34</sup>CHILIKIN 19 reports signals in  $B^+ \rightarrow \eta_c(2S)K^+$  and  $B^0 \rightarrow \eta_c(2S)K_S^0$  with 12.3 and 5.9 standard deviations, respectively.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4 \times 10^{-4}$	90	<sup>35</sup> WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
$<0.01$	90	LEE	85	CBAL $\psi' \rightarrow \text{photons}$

<sup>35</sup> WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S) K^+) = 4.4 \times 10^{-4}$ .

$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi)$   $\Gamma_{20}/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt;3.33</math></b>	90	<sup>36</sup> LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$
------------------------------	----	--------------------	-----------	---

<sup>36</sup> We divided the reported limit by 3 to take into account isospin relations.

**$\eta_c(2S)$  CROSS-PARTICLE BRANCHING RATIOS**

$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<11.8 \times 10^{-6}$	90	<sup>37</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
------------------------	----	------------------------------	------	--

<sup>37</sup> CRONIN-HENNESSY 10 reports a limit of  $< 5.9 \times 10^{-6}$  for the decay  $\eta_c(2S) \rightarrow K^+ K^- \eta$  which we multiply by 2 account for isospin symmetry. It assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. It also gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt;14.6 \times 10^{-6}</math></b>	90	<sup>38</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
---	----	------------------------------	------	---

<sup>38</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt;12.7 \times 10^{-7}</math></b>	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
---	----	---------	----------	---

$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt;13.2 \times 10^{-6}</math></b>	90	<sup>39</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
---	----	------------------------------	------	---

<sup>39</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_7 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	40 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>40</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_8 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	41 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>41</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{10} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	42 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>42</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{11} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM 13K	BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15.2$	90	43 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

<sup>43</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{13} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM 13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33	44 ABLIKIM 17N	BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$

<sup>44</sup> Uses  $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$ .

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	45 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

<sup>45</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	46 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>46</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	47 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$

<sup>47</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

## $\eta_c(2S)$ REFERENCES

CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)

UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

---