

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on “ $\psi(2S)$  and  $\chi_c$  branching ratios” before the  $\chi_{c0}(1P)$  Listings.

### $\psi(2S)$ MASS

OUR FIT includes measurements of  $m_{\psi(2S)}$ ,  $m_{\psi(3770)}$ , and  $m_{\psi(3770)} - m_{\psi(2S)}$ .

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3686.10 ± 0.06 OUR FIT</b>				Error includes scale factor of 5.9.
<b>3686.097 ± 0.010 OUR AVERAGE</b>				
3686.099 ± 0.004 ± 0.009		<sup>1</sup> ANASHIN 15	KEDR	$e^+e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ 12H	LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	<sup>2</sup> ARTAMONOV 00	OLYA	$e^+e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		<sup>3</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3686.114 ± 0.007 <sup>+0.011</sup> <sub>-0.016</sub>		<sup>4</sup> ANASHIN 12	KEDR	$e^+e^- \rightarrow$ hadrons
3686.111 ± 0.025 ± 0.009		AULCHENKO 03	KEDR	$e^+e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	<sup>5</sup> ZHOLENTZ 80	OLYA	$e^+e^-$

<sup>1</sup> Supersedes AULCHENKO 03 and ANASHIN 12.

<sup>2</sup> Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

<sup>3</sup> Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the  $J/\psi(1S)$  mass from AULCHENKO 03.

<sup>4</sup> From the scans in 2004 and 2006. ANASHIN 12 reports the value  $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$  MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

<sup>5</sup> Superseded by ARTAMONOV 00.

### $m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>589.188 ± 0.028 OUR AVERAGE</b>			
589.194 ± 0.027 ± 0.011	<sup>1</sup> AULCHENKO 03	KEDR	$e^+e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	<sup>1</sup> ZHOLENTZ 80	OLYA	$e^+e^-$
588.7 ± 0.8	LUTH 75	MRK1	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
588 ± 1	<sup>2</sup> BAI 98E	BES	$e^+e^-$

<sup>1</sup> Redundant with data in mass above.

<sup>2</sup> Systematic errors not evaluated.

**$\psi(2S)$  WIDTH**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>294 ± 8 OUR FIT</b>				
<b>286 ± 16 OUR AVERAGE</b>				
358 ± 88 ± 4		ABLIKIM	08B BES2	$e^+e^- \rightarrow \text{hadrons}$
290 ± 25 ± 4	2.7k	ANDREOTTI	07 E835	$\rho\bar{p} \rightarrow e^+e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM	06L BES2	$e^+e^- \rightarrow \text{hadrons}$
264 ± 27		<sup>1</sup> BAI	02B BES2	$e^+e^-$
287 ± 37 ± 16		<sup>2</sup> ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$

<sup>1</sup>From a simultaneous fit to the hadronic and  $\mu^+\mu^-$  cross section, assuming  $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$  and lepton universality. Does not include vacuum polarization correction.

<sup>2</sup>The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

 **$\psi(2S)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(97.85 ± 0.13) %	
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	( 1.73 ± 0.14 ) %	S=1.5
$\Gamma_3$ $ggg$	(10.6 ± 1.6) %	
$\Gamma_4$ $\gamma gg$	( 1.03 ± 0.29 ) %	
$\Gamma_5$ light hadrons	(15.4 ± 1.5) %	
$\Gamma_6$ $K_S^0$ anything	(16.0 ± 1.1) %	
$\Gamma_7$ $e^+e^-$	( 7.93 ± 0.17 ) × 10 <sup>-3</sup>	
$\Gamma_8$ $\mu^+\mu^-$	( 8.0 ± 0.6 ) × 10 <sup>-3</sup>	
$\Gamma_9$ $\tau^+\tau^-$	( 3.1 ± 0.4 ) × 10 <sup>-3</sup>	

**Decays into  $J/\psi(1S)$  and anything**

$\Gamma_{10}$ $J/\psi(1S)$ anything	(61.4 ± 0.6) %	
$\Gamma_{11}$ $J/\psi(1S)$ neutrals	(25.38 ± 0.32) %	
$\Gamma_{12}$ $J/\psi(1S)\pi^+\pi^-$	(34.68 ± 0.30) %	
$\Gamma_{13}$ $J/\psi(1S)\pi^0\pi^0$	(18.24 ± 0.31) %	
$\Gamma_{14}$ $J/\psi(1S)\eta$	( 3.37 ± 0.05 ) %	
$\Gamma_{15}$ $J/\psi(1S)\pi^0$	( 1.268 ± 0.032 ) × 10 <sup>-3</sup>	

**Hadronic decays**

$\Gamma_{16}$ $\pi^+\pi^-$	( 7.8 ± 2.6 ) × 10 <sup>-6</sup>	
$\Gamma_{17}$ $\pi^+\pi^-\pi^0$	( 2.01 ± 0.17 ) × 10 <sup>-4</sup>	S=1.7
$\Gamma_{18}$ $\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$	( 3.2 ± 1.2 ) × 10 <sup>-5</sup>	S=1.8
$\Gamma_{19}$ $\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	( 1.9 <sup>+1.2</sup> / <sub>-0.4</sub> ) × 10 <sup>-4</sup>	
$\Gamma_{20}$ $2(\pi^+\pi^-)$	( 2.4 ± 0.6 ) × 10 <sup>-4</sup>	S=2.2
$\Gamma_{21}$ $\rho^0\pi^+\pi^-$	( 2.2 ± 0.6 ) × 10 <sup>-4</sup>	S=1.4
$\Gamma_{22}$ $2(\pi^+\pi^-)\pi^0$	( 2.9 ± 1.0 ) × 10 <sup>-3</sup>	S=4.7
$\Gamma_{23}$ $\rho a_2(1320)$	( 2.6 ± 0.9 ) × 10 <sup>-4</sup>	
$\Gamma_{24}$ $\pi^+\pi^-\pi^0\pi^0\pi^0$	( 5.3 ± 0.9 ) × 10 <sup>-3</sup>	

$\Gamma_{25}$	$\pi^+\pi^-4\pi^0$	$(1.4 \pm 1.0) \times 10^{-3}$	
$\Gamma_{26}$	$\rho^\pm\pi^\mp\pi^0\pi^0$	$< 2.7 \times 10^{-3}$	CL=90%
$\Gamma_{27}$	$3(\pi^+\pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8
$\Gamma_{28}$	$2(\pi^+\pi^-\pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$	
$\Gamma_{29}$	$3(\pi^+\pi^-\pi^0)$	$(3.5 \pm 1.6) \times 10^{-3}$	
$\Gamma_{30}$	$2(\pi^+\pi^-)3\pi^0$	$(1.42 \pm 0.31) \%$	
$\Gamma_{31}$	$\eta\pi^+\pi^-$	$< 1.6 \times 10^{-4}$	CL=90%
$\Gamma_{32}$	$\eta\pi^+\pi^-\pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$	
$\Gamma_{33}$	$\eta 2(\pi^+\pi^-)$	$(1.2 \pm 0.6) \times 10^{-3}$	
$\Gamma_{34}$	$\eta\pi^+\pi^-\pi^0\pi^0$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{35}$	$\eta\pi^+\pi^-3\pi^0$	$< 2.1 \times 10^{-3}$	CL=90%
$\Gamma_{36}$	$\eta 2(\pi^+\pi^-\pi^0)$	$< 2.1 \times 10^{-3}$	CL=90%
$\Gamma_{37}$	$\rho\eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1
$\Gamma_{38}$	$\eta'\pi^+\pi^-\pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$	
$\Gamma_{39}$	$\eta'\rho$	$(1.9 \pm 1.7 \mp 1.2) \times 10^{-5}$	
$\Gamma_{40}$	$\omega\pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$	
$\Gamma_{41}$	$\omega\pi^+\pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1
$\Gamma_{42}$	$\omega\pi^+\pi^-2\pi^0$	$(8.7 \pm 2.4) \times 10^{-3}$	
$\Gamma_{43}$	$b_1^\pm\pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1
$\Gamma_{44}$	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$	
$\Gamma_{45}$	$\omega\pi^0\pi^0$	$(1.11 \pm 0.35) \times 10^{-3}$	
$\Gamma_{46}$	$\omega 3\pi^0$	$< 8 \times 10^{-4}$	CL=90%
$\Gamma_{47}$	$b_1^0\pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$	
$\Gamma_{48}$	$\omega\eta$	$< 1.1 \times 10^{-5}$	CL=90%
$\Gamma_{49}$	$\omega\eta'$	$(3.2 \pm 2.5 \mp 2.1) \times 10^{-5}$	
$\Gamma_{50}$	$\phi\pi^0$	$< 4 \times 10^{-7}$	CL=90%
$\Gamma_{51}$	$\phi\pi^+\pi^-$	$(1.18 \pm 0.26) \times 10^{-4}$	S=1.5
$\Gamma_{52}$	$\phi f_0(980) \rightarrow \pi^+\pi^-$	$(7.5 \pm 3.3) \times 10^{-5}$	S=1.6
$\Gamma_{53}$	$\phi\eta$	$(3.10 \pm 0.31) \times 10^{-5}$	
$\Gamma_{54}$	$\eta\phi(2170), \phi(2170) \rightarrow$ $\phi f_0(980), f_0 \rightarrow \pi^+\pi^-$	$< 2.2 \times 10^{-6}$	CL=90%
$\Gamma_{55}$	$\phi\eta'$	$(1.54 \pm 0.20) \times 10^{-5}$	
$\Gamma_{56}$	$\phi f_1(1285)$	$(3.0 \pm 1.3) \times 10^{-5}$	
$\Gamma_{57}$	$\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta$	$(8.5 \pm 1.7) \times 10^{-6}$	
$\Gamma_{58}$	$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$	
$\Gamma_{59}$	$K^+K^-$	$(7.5 \pm 0.5) \times 10^{-5}$	
$\Gamma_{60}$	$K^+K^-\pi^+$	$(7.3 \pm 0.5) \times 10^{-4}$	
$\Gamma_{61}$	$K^+K^-\pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$	
$\Gamma_{62}$	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$	
$\Gamma_{63}$	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$	
$\Gamma_{64}$	$K_S^0 K_L^0\pi^0$	$< 3.0 \times 10^{-4}$	CL=90%
$\Gamma_{65}$	$K^+K^-\pi^0\pi^0$	$(2.6 \pm 1.3) \times 10^{-4}$	

$\Gamma_{66}$	$K^+ K^- \pi^+ \pi^- \pi^0$	$( 1.26 \pm 0.09 ) \times 10^{-3}$	
$\Gamma_{67}$	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$( 5.9 \pm 2.2 ) \times 10^{-5}$	
$\Gamma_{68}$	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$( 8.6 \pm 2.2 ) \times 10^{-4}$	
$\Gamma_{69}$	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$( 9.6 \pm 2.8 ) \times 10^{-4}$	
$\Gamma_{70}$	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$( 7.3 \pm 2.6 ) \times 10^{-4}$	
$\Gamma_{71}$	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$( 6.1 \pm 1.8 ) \times 10^{-4}$	
$\Gamma_{72}$	$K_S^0 K_S^0 \pi^+ \pi^-$	$( 2.2 \pm 0.4 ) \times 10^{-4}$	
$\Gamma_{73}$	$K_S^0 K_L^0 \pi^0 \pi^0$	$( 1.3 \pm 0.6 ) \times 10^{-3}$	
$\Gamma_{74}$	$K_S^0 K_L^0 \eta$	$( 1.3 \pm 0.5 ) \times 10^{-3}$	
$\Gamma_{75}$	$K^+ K^- \rho^0$	$( 2.2 \pm 0.4 ) \times 10^{-4}$	
$\Gamma_{76}$	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$( 1.9 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{77}$	$K^+ K^- \pi^+ \pi^- \eta$	$( 1.3 \pm 0.7 ) \times 10^{-3}$	
$\Gamma_{78}$	$K^+ K^- 2(\pi^+ \pi^-)$	$( 1.9 \pm 0.9 ) \times 10^{-3}$	
$\Gamma_{79}$	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$( 1.00 \pm 0.31 ) \times 10^{-3}$	
$\Gamma_{80}$	$K^+ K^*(892)^- + \text{c.c.}$	$( 2.9 \pm 0.4 ) \times 10^{-5}$	S=1.2
$\Gamma_{81}$	$2(K^+ K^-)$	$( 6.3 \pm 1.3 ) \times 10^{-5}$	
$\Gamma_{82}$	$2(K^+ K^-) \pi^0$	$( 1.10 \pm 0.28 ) \times 10^{-4}$	
$\Gamma_{83}$	$K^+ K^- \phi$	$( 7.0 \pm 1.6 ) \times 10^{-5}$	
$\Gamma_{84}$	$K_1(1270)^\pm K^\mp$	$( 1.00 \pm 0.28 ) \times 10^{-3}$	
$\Gamma_{85}$	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$( 6.7 \pm 2.5 ) \times 10^{-4}$	
$\Gamma_{86}$	$\eta K^+ K^-$ , no $\eta \phi$	$( 3.49 \pm 0.17 ) \times 10^{-5}$	
$\Gamma_{87}$	$X(1750) \eta \rightarrow K^+ K^- \eta$	$( 4.8 \pm 2.8 ) \times 10^{-6}$	
$\Gamma_{88}$	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%
$\Gamma_{89}$	$K_2^*(1430)^\pm K^\mp$	$( 7.1 \begin{smallmatrix} +1.3 \\ -0.9 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{90}$	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$( 1.09 \pm 0.20 ) \times 10^{-4}$	
$\Gamma_{91}$	$\omega K^+ K^-$	$( 1.62 \pm 0.11 ) \times 10^{-4}$	S=1.1
$\Gamma_{92}$	$\omega K_S^0 K_S^0$	$( 7.0 \pm 0.5 ) \times 10^{-5}$	
$\Gamma_{93}$	$\omega K^*(892)^+ K^- + \text{c.c.}$	$( 2.07 \pm 0.26 ) \times 10^{-4}$	
$\Gamma_{94}$	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$( 6.1 \pm 1.2 ) \times 10^{-5}$	
$\Gamma_{95}$	$\omega \bar{K}^*(892)^0 K^0$	$( 1.68 \pm 0.30 ) \times 10^{-4}$	
$\Gamma_{96}$	$\omega \bar{K}_2^*(1430)^0 K^0$	$( 5.8 \pm 2.2 ) \times 10^{-5}$	
$\Gamma_{97}$	$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$( 1.6 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{98}$	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$( 1.09 \pm 0.26 ) \times 10^{-5}$	
$\Gamma_{99}$	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$( 3.0 \pm 1.0 ) \times 10^{-6}$	
$\Gamma_{100}$	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$( 1.2 \pm 0.7 ) \times 10^{-6}$	
$\Gamma_{101}$	$p \bar{p}$	$( 2.94 \pm 0.08 ) \times 10^{-4}$	
$\Gamma_{102}$	$n \bar{n}$	$( 3.06 \pm 0.15 ) \times 10^{-4}$	
$\Gamma_{103}$	$p \bar{p} \pi^0$	$( 1.53 \pm 0.07 ) \times 10^{-4}$	
$\Gamma_{104}$	$N(940) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$	$( 6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{105}$	$N(1440) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$	$( 7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix} ) \times 10^{-5}$	S=2.5

$\Gamma_{106}$	$N(1520)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix} ) \times 10^{-6}$	
$\Gamma_{107}$	$N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 2.5 \pm 1.0 ) \times 10^{-5}$	
$\Gamma_{108}$	$N(1650)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{109}$	$N(1720)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{110}$	$N(2300)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{111}$	$N(2570)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0$	$( 2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{112}$	$p\bar{p}\pi^+\pi^-$	$( 6.0 \pm 0.4 ) \times 10^{-4}$	
$\Gamma_{113}$	$p\bar{p}K^+K^-$	$( 2.7 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{114}$	$p\bar{p}\eta$	$( 6.0 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{115}$	$N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\eta$	$( 4.5 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{116}$	$p\bar{p}\pi^+\pi^-\pi^0$	$( 7.3 \pm 0.7 ) \times 10^{-4}$	
$\Gamma_{117}$	$p\bar{p}\rho^0$	$( 5.0 \pm 2.2 ) \times 10^{-5}$	
$\Gamma_{118}$	$p\bar{p}\omega$	$( 6.9 \pm 2.1 ) \times 10^{-5}$	
$\Gamma_{119}$	$p\bar{p}\eta'$	$( 1.10 \pm 0.13 ) \times 10^{-5}$	
$\Gamma_{120}$	$p\bar{p}\phi$	$( 6.1 \pm 0.6 ) \times 10^{-6}$	
$\Gamma_{121}$	$\phi X(1835) \rightarrow p\bar{p}\phi$	$< 1.82 \times 10^{-7}$	CL=90%
$\Gamma_{122}$	$p\bar{n}\pi^-\text{ or c.c.}$	$( 2.48 \pm 0.17 ) \times 10^{-4}$	
$\Gamma_{123}$	$p\bar{n}\pi^-\pi^0$	$( 3.2 \pm 0.7 ) \times 10^{-4}$	
$\Gamma_{124}$	$\Lambda\bar{\Lambda}$	$( 3.81 \pm 0.13 ) \times 10^{-4}$	S=1.4
$\Gamma_{125}$	$\Lambda\bar{\Lambda}\pi^0$	$< 2.9 \times 10^{-6}$	CL=90%
$\Gamma_{126}$	$\Lambda\bar{\Lambda}\eta$	$( 2.5 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{127}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$( 2.8 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{128}$	$\Lambda\bar{p}K^+$	$( 1.00 \pm 0.14 ) \times 10^{-4}$	
$\Gamma_{129}$	$\Lambda\bar{p}K^*(892)^+ + \text{c.c.}$	$( 6.3 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{130}$	$\Lambda\bar{p}K^+\pi^+\pi^-$	$( 1.8 \pm 0.4 ) \times 10^{-4}$	
$\Gamma_{131}$	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$( 8.1 \pm 1.8 ) \times 10^{-5}$	
$\Gamma_{132}$	$\Delta^{++}\bar{\Delta}^{--}$	$( 1.28 \pm 0.35 ) \times 10^{-4}$	
$\Gamma_{133}$	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$( 1.40 \pm 0.13 ) \times 10^{-4}$	
$\Gamma_{134}$	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	$( 1.54 \pm 0.14 ) \times 10^{-4}$	
$\Gamma_{135}$	$\Lambda\bar{\Sigma}^0 + \text{c.c.}$	$( 1.6 \pm 0.7 ) \times 10^{-6}$	
$\Gamma_{136}$	$\Lambda\bar{\Sigma}^0$		
$\Gamma_{137}$	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	$( 1.67 \pm 0.18 ) \times 10^{-5}$	
$\Gamma_{138}$	$\Sigma^+\bar{\Sigma}^-$	$( 2.43 \pm 0.10 ) \times 10^{-4}$	S=1.4
$\Gamma_{139}$	$\Sigma^0\bar{\Sigma}^0$	$( 2.35 \pm 0.09 ) \times 10^{-4}$	S=1.1
$\Gamma_{140}$	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$( 8.5 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{141}$	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$( 8.5 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{142}$	$\Sigma(1385)^0\bar{\Sigma}(1385)^0$	$( 6.9 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{143}$	$\Xi^-\bar{\Xi}^+$	$( 2.87 \pm 0.11 ) \times 10^{-4}$	S=1.1
$\Gamma_{144}$	$\Xi^0\bar{\Xi}^0$	$( 2.3 \pm 0.4 ) \times 10^{-4}$	S=4.2
$\Gamma_{145}$	$\Xi(1530)^0\bar{\Xi}(1530)^0$	$( 6.8 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{146}$	$\Lambda\bar{\Xi}^+K^- + \text{c.c.}$	$( 3.9 \pm 0.4 ) \times 10^{-5}$	

$\Gamma_{147}$	$\Xi(1530)^- \Xi(1530)^+$	$( 1.15 \pm 0.07 ) \times 10^{-4}$	
$\Gamma_{148}$	$\Xi(1530)^- \Xi^+$	$( 7.0 \pm 1.2 ) \times 10^{-6}$	
$\Gamma_{149}$	$\Xi(1530)^0 \Xi^0$	$( 5.3 \pm 0.5 ) \times 10^{-6}$	
$\Gamma_{150}$	$\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ +$	$( 5.2 \pm 1.6 ) \times 10^{-6}$	
$\Gamma_{151}$	$\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ +$	$( 1.20 \pm 0.32 ) \times 10^{-5}$	
$\Gamma_{152}$	$\Sigma^0 \Xi^+ K^- + \text{c.c.}$	$( 3.7 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{153}$	$\Omega^- \bar{\Omega}^+$	$( 5.66 \pm 0.30 ) \times 10^{-5}$	S=1.3
$\Gamma_{154}$	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%
$\Gamma_{155}$	$h_c(1P) \pi^0$	$( 8.6 \pm 1.3 ) \times 10^{-4}$	
$\Gamma_{156}$	$\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$	$< 1.7 \times 10^{-6}$	CL=90%
$\Gamma_{157}$	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	[a] $< 8.8 \times 10^{-6}$	CL=90%
$\Gamma_{158}$	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 1.0 \times 10^{-5}$	CL=90%
$\Gamma_{159}$	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 7.0 \times 10^{-6}$	CL=90%
$\Gamma_{160}$	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 2.6 \times 10^{-5}$	CL=90%
$\Gamma_{161}$	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 6.0 \times 10^{-6}$	CL=90%

### Radiative decays

$\Gamma_{162}$	$\gamma \chi_{c0}(1P)$	$( 9.79 \pm 0.20 ) \%$	
$\Gamma_{163}$	$\gamma \chi_{c1}(1P)$	$( 9.75 \pm 0.24 ) \%$	
$\Gamma_{164}$	$\gamma \chi_{c2}(1P)$	$( 9.52 \pm 0.20 ) \%$	
$\Gamma_{165}$	$\gamma \eta_c(1S)$	$( 3.4 \pm 0.5 ) \times 10^{-3}$	S=1.3
$\Gamma_{166}$	$\gamma \eta_c(2S)$	$( 7 \pm 5 ) \times 10^{-4}$	
$\Gamma_{167}$	$\gamma \pi^0$	$( 1.04 \pm 0.22 ) \times 10^{-6}$	S=1.4
$\Gamma_{168}$	$\gamma 2(\pi^+ \pi^-)$	$( 4.0 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{169}$	$\gamma 3(\pi^+ \pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%
$\Gamma_{170}$	$\gamma \eta'(958)$	$( 1.24 \pm 0.04 ) \times 10^{-4}$	
$\Gamma_{171}$	$\gamma f_2(1270)$	$( 2.73^{+0.29}_{-0.25} ) \times 10^{-4}$	S=1.8
$\Gamma_{172}$	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$( 3.1 \pm 1.7 ) \times 10^{-5}$	
$\Gamma_{173}$	$\gamma f_0(1500)$	$( 9.3 \pm 1.9 ) \times 10^{-5}$	
$\Gamma_{174}$	$\gamma f_2'(1525)$	$( 3.3 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{175}$	$\gamma f_0(1710)$		
$\Gamma_{176}$	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$( 3.5 \pm 0.6 ) \times 10^{-5}$	
$\Gamma_{177}$	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$( 6.6 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{178}$	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$( 4.8 \pm 1.0 ) \times 10^{-6}$	
$\Gamma_{179}$	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$( 3.2 \pm 1.0 ) \times 10^{-6}$	
$\Gamma_{180}$	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 5.8 \times 10^{-6}$	CL=90%
$\Gamma_{181}$	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%
$\Gamma_{182}$	$\gamma \eta$	$( 9.2 \pm 1.8 ) \times 10^{-7}$	
$\Gamma_{183}$	$\gamma \eta \pi^+ \pi^-$	$( 8.7 \pm 2.1 ) \times 10^{-4}$	
$\Gamma_{184}$	$\gamma \eta(1405)$		
$\Gamma_{185}$	$\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$	$< 9 \times 10^{-5}$	CL=90%

$\Gamma_{186}$	$\gamma\eta(1405) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$	
$\Gamma_{187}$	$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0$	$< 5.0 \times 10^{-7}$	CL=90%
$\Gamma_{188}$	$\gamma\eta(1475)$		
$\Gamma_{189}$	$\gamma\eta(1475) \rightarrow \gamma K\bar{K}\pi$	$< 1.4 \times 10^{-4}$	CL=90%
$\Gamma_{190}$	$\gamma\eta(1475) \rightarrow \gamma\eta\pi^+\pi^-$	$< 8.8 \times 10^{-5}$	CL=90%
$\Gamma_{191}$	$\gamma K^{*0}K^+\pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$	
$\Gamma_{192}$	$\gamma K^{*0}\bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$	
$\Gamma_{193}$	$\gamma K_S^0 K^+\pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$	
$\Gamma_{194}$	$\gamma K^+ K^-\pi^+\pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	
$\Gamma_{195}$	$\gamma K^+ K^- 2(\pi^+\pi^-)$	$< 2.2 \times 10^{-4}$	CL=90%
$\Gamma_{196}$	$\gamma 2(K^+K^-)$	$< 4 \times 10^{-5}$	CL=90%
$\Gamma_{197}$	$\gamma p\bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0
$\Gamma_{198}$	$\gamma f_2(1950) \rightarrow \gamma p\bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$	
$\Gamma_{199}$	$\gamma f_2(2150) \rightarrow \gamma p\bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$	
$\Gamma_{200}$	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	$(4.6 \pm_{-4.0}^{+1.8}) \times 10^{-6}$	
$\Gamma_{201}$	$\gamma X \rightarrow \gamma p\bar{p}$	[b] $< 2 \times 10^{-6}$	CL=90%
$\Gamma_{202}$	$\gamma p\bar{p}\pi^+\pi^-$	$(2.8 \pm 1.4) \times 10^{-5}$	
$\Gamma_{203}$	$\gamma\gamma$	$< 1.5 \times 10^{-4}$	CL=90%
$\Gamma_{204}$	$\gamma\gamma J/\psi$	$(3.1 \pm_{-1.2}^{+1.0}) \times 10^{-4}$	
$\Gamma_{205}$	$e^+e^-\eta'$	$(1.90 \pm 0.26) \times 10^{-6}$	
$\Gamma_{206}$	$e^+e^-\chi_{c0}(1P)$	$(1.06 \pm 0.24) \times 10^{-3}$	
$\Gamma_{207}$	$e^+e^-\chi_{c1}(1P)$	$(8.5 \pm 0.6) \times 10^{-4}$	
$\Gamma_{208}$	$e^+e^-\chi_{c2}(1P)$	$(7.0 \pm 0.8) \times 10^{-4}$	

#### Weak decays

$\Gamma_{209}$	$D^0 e^+e^- + \text{c.c.}$	$< 1.4 \times 10^{-7}$	CL=90%
----------------	----------------------------	------------------------	--------

#### Other decays

$\Gamma_{210}$	invisible	$< 1.6 \%$	CL=90%
----------------	-----------	------------	--------

[a]  $\Theta(1540)$  is a hypothetical pentaquark state of 1.54 GeV/c<sup>2</sup> mass and a width of less than 25 MeV/c<sup>2</sup>.

[b] For a narrow resonance in the range  $2.2 < M(X) < 2.8$  GeV.

---

### CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

x <sub>8</sub>	3									
x <sub>9</sub>	1	0								
x <sub>12</sub>	29	11	2							
x <sub>13</sub>	28	6	1	48						
x <sub>14</sub>	13	4	1	36	15					
x <sub>101</sub>	0	0	0	4	3	2				
x <sub>162</sub>	1	0	0	2	1	1	0			
x <sub>163</sub>	1	0	0	2	1	1	0	0		
x <sub>164</sub>	1	0	0	3	1	1	0	0	0	
Γ	-81	-4	-1	-38	-34	-16	-7	-1	-1	-1
	x <sub>7</sub>	x <sub>8</sub>	x <sub>9</sub>	x <sub>12</sub>	x <sub>13</sub>	x <sub>14</sub>	x <sub>101</sub>	x <sub>162</sub>	x <sub>163</sub>	x <sub>164</sub>

### ψ(2S) PARTIAL WIDTHS

#### Γ(hadrons)

Γ<sub>1</sub>

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
258 ± 26	BAI	02B	BES2 e <sup>+</sup> e <sup>-</sup>
224 ± 56	LUTH	75	MRK1 e <sup>+</sup> e <sup>-</sup>

#### Γ(e<sup>+</sup>e<sup>-</sup>)

Γ<sub>7</sub>

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>2.33 ± 0.04 OUR FIT</b>			
<b>2.29 ± 0.06 OUR AVERAGE</b>			
2.23 ± 0.10 ± 0.02	<sup>1</sup> ABLIKIM	15V	BES3 4.0–4.4 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> J/ψ
2.338 ± 0.037 ± 0.096	ABLIKIM	08B	BES2 e <sup>+</sup> e <sup>-</sup> → hadrons
2.330 ± 0.036 ± 0.110	ABLIKIM	06L	BES2 e <sup>+</sup> e <sup>-</sup> → hadrons
2.44 ± 0.21	<sup>2</sup> BAI	02B	BES2 e <sup>+</sup> e <sup>-</sup>
2.14 ± 0.21	ALEXANDER	89	RVUE See γ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2.279 ± 0.015 ± 0.042	<sup>3</sup> ANASHIN	18	KEDR e <sup>+</sup> e <sup>-</sup>
2.282 ± 0.015 ± 0.042	<sup>4</sup> ANASHIN	18	KEDR e <sup>+</sup> e <sup>-</sup>
2.0 ± 0.3	BRANDELIK	79C	DASP e <sup>+</sup> e <sup>-</sup>
2.1 ± 0.3	<sup>5</sup> LUTH	75	MRK1 e <sup>+</sup> e <sup>-</sup>

<sup>1</sup> ABLIKIM 15V reports 2.213 ± 0.018 ± 0.099 keV from a measurement of [Γ(ψ(2S) → e<sup>+</sup>e<sup>-</sup>)] × [B(ψ(2S) → J/ψ(1S)π<sup>+</sup>π<sup>-</sup>)] assuming B(ψ(2S) → J/ψ(1S)π<sup>+</sup>π<sup>-</sup>) = (34.95 ± 0.45) × 10<sup>-2</sup>, which we rescale to our best value B(ψ(2S) → J/ψ(1S)π<sup>+</sup>π<sup>-</sup>) = (34.68 ± 0.30) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> From a simultaneous fit to e<sup>+</sup>e<sup>-</sup>, μ<sup>+</sup>μ<sup>-</sup>, and hadronic channel, assuming Γ<sub>e</sub> = Γ<sub>μ</sub> = Γ<sub>τ</sub>/0.38847.

<sup>3</sup> Combining Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(μ<sup>+</sup>μ<sup>-</sup>) from ANASHIN 18 with Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(hadrons) from ANASHIN 12 and assuming lepton universality.

<sup>4</sup> From the sum of Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(hadrons) from ANASHIN 12, Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(e<sup>+</sup>e<sup>-</sup>) and Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(μ<sup>+</sup>μ<sup>-</sup>) from ANASHIN 18, and Γ<sub>e<sup>+</sup>e<sup>-</sup></sub> · B(τ<sup>+</sup>τ<sup>-</sup>) from ANASHIN 07.

<sup>5</sup> From a simultaneous fit to e<sup>+</sup>e<sup>-</sup>, μ<sup>+</sup>μ<sup>-</sup>, and hadronic channels assuming Γ(e<sup>+</sup>e<sup>-</sup>) = Γ(μ<sup>+</sup>μ<sup>-</sup>).



$\Gamma(\gamma\gamma)$					$\Gamma_{203}$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<43	90	BRANDELIK	79C	DASP	$e^+e^-$

### $\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the integrated cross section into channel(i) in the  $e^+e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(i)$  or the branching ratio  $\Gamma(i)/\text{total}$ .

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_7/\Gamma$
VALUE (keV)		DOCUMENT ID	TECN	COMMENT	
<b>2.233±0.015±0.042</b>		<sup>1</sup> ANASHIN	12	KEDR	$e^+e^- \rightarrow \text{hadrons}$
• • •					We do not use the following data for averages, fits, limits, etc. • • •
2.2 ±0.4		ABRAMS	75	MRK1	$e^+e^-$

<sup>1</sup> ANASHIN 12 reports the value  $2.233 \pm 0.015 \pm 0.037 \pm 0.020$  keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

$\Gamma(K_S^0 \text{ anything}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_6\Gamma_7/\Gamma$
VALUE (keV)		DOCUMENT ID	TECN	COMMENT	
<b>0.3738±0.0067±0.0200</b>		ABLIKIM	21S	BES3	$e^+e^- \rightarrow K_S^0 \text{ anything}$

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_7/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
<b>21.2±0.7±1.2</b>		<sup>1</sup> ANASHIN	18	KEDR	$e^+e^-$
					<sup>1</sup> From the average of nine scans of the $\psi(2S)$ .


$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_7/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
<b>19.3±0.3±0.5</b>		<sup>1</sup> ANASHIN	18	KEDR	$\psi(2S) \rightarrow \mu^+\mu^-$
					<sup>1</sup> From the average of nine scans of the $\psi(2S)$ .

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_9\Gamma_7/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • •					We do not use the following data for averages, fits, limits, etc. • • •
9.0±2.6	79	<sup>1</sup> ANASHIN	07	KEDR	$e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$
					<sup>1</sup> Using $\psi(2S)$ total width of $337 \pm 13$ keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{12}\Gamma_7/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.808±0.013 OUR FIT</b>					
<b>0.837±0.025 OUR AVERAGE</b>					Error includes scale factor of 1.3. See the ideogram below.
0.837±0.028±0.005		<sup>1</sup> LEES	12E	BABR	10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$
0.852±0.010±0.026	19.5k	ADAM	06	CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$
0.68 ±0.09		<sup>2</sup> BAI	98E	BES	$e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.03    256    <sup>3</sup> AUBERT 07AU BABR 10.6 e<sup>+</sup>e<sup>-</sup> → J/ψ π<sup>+</sup>π<sup>-</sup>γ  
 0.755 ± 0.048 ± 0.004    544    <sup>4</sup> AUBERT 05D BABR 10.6 e<sup>+</sup>e<sup>-</sup> → π<sup>+</sup>π<sup>-</sup>μ<sup>+</sup>μ<sup>-</sup>γ



<sup>1</sup> LEES 12E reports [ $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(J/ψ(1S) → μ<sup>+</sup>μ<sup>-</sup>)] = (49.9 ± 1.3 ± 1.0) × 10<sup>-3</sup> keV which we divide by our best value B(J/ψ(1S) → μ<sup>+</sup>μ<sup>-</sup>) = (5.961 ± 0.033) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> The value of  $\Gamma(e^+e^-)$  quoted in BAI 98E is derived using B(ψ(2S) → J/ψ(1S)π<sup>+</sup>π<sup>-</sup>) = (32.4 ± 2.6) × 10<sup>-2</sup> and B(J/ψ(1S) → ℓ<sup>+</sup>ℓ<sup>-</sup>) = 0.1203 ± 0.0038. Recalculated by us using B(J/ψ(1S) → ℓ<sup>+</sup>ℓ<sup>-</sup>) = 0.1181 ± 0.0020.

<sup>3</sup> AUBERT 07AU reports [ $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(J/ψ(1S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>)] = 0.0186 ± 0.0012 ± 0.0011 keV which we divide by our best value B(J/ψ(1S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) = (2.10 ± 0.08) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> AUBERT 05D reports [ $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(J/ψ(1S) → μ<sup>+</sup>μ<sup>-</sup>)] = 0.0450 ± 0.0018 ± 0.0022 keV which we divide by our best value B(J/ψ(1S) → μ<sup>+</sup>μ<sup>-</sup>) = (5.961 ± 0.033) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  (keV)

$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{13}\Gamma_7/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.425 ± 0.009 OUR FIT</b>					
<b>0.411 ± 0.008 ± 0.018</b>	3.6k	ADAM	06 CLEO	3.773 e <sup>+</sup> e <sup>-</sup> → γψ(2S)	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48 ± 0.09 ± 0.02      142    <sup>1</sup> LEES      18E BABR    10.6 e<sup>+</sup>e<sup>-</sup> → J/ψ π<sup>0</sup> π<sup>0</sup> γ

<sup>1</sup> LEES 18E reports [Γ(ψ(2S) → J/ψ(1S)π<sup>0</sup>π<sup>0</sup>) × Γ(ψ(2S) → e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>] × [B(J/ψ(1S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>)] = 0.0101 ± 0.0015 ± 0.0011 keV which we divide by our best value B(J/ψ(1S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) = (2.10 ± 0.08) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**Γ(J/ψ(1S)η) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>14</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**78.6 ± 1.6 OUR FIT**

**87 ± 9 OUR AVERAGE**

83 ± 25 ± 5      14      <sup>1</sup> AUBERT      07AU BABR    10.6 e<sup>+</sup>e<sup>-</sup> → J/ψ π<sup>+</sup>π<sup>-</sup>π<sup>0</sup> γ

88 ± 6 ± 7      291 ± 24      ADAM      06      CLEO    3.773 e<sup>+</sup>e<sup>-</sup> → γψ(2S)

<sup>1</sup> AUBERT 07AU quotes Γ<sub>ee</sub><sup>ψ(2S)</sup> · B(ψ(2S) → J/ψη) · B(J/ψ → μ<sup>+</sup>μ<sup>-</sup>) · B(η → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) = 1.11 ± 0.33 ± 0.07 eV.

**Γ(J/ψ(1S)π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>15</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      CL%      EVTS      DOCUMENT ID      TECN      COMMENT

**<8**      90      <37      ADAM      06      CLEO    3.773 e<sup>+</sup>e<sup>-</sup> → γψ(2S)

**Γ(2(π<sup>+</sup>π<sup>-</sup>)π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>22</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**29.7 ± 2.2 ± 1.8**      410      AUBERT      07AU BABR    10.6 e<sup>+</sup>e<sup>-</sup> → 2(π<sup>+</sup>π<sup>-</sup>)π<sup>0</sup> γ

**Γ(π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>π<sup>0</sup>π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>24</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**12.4 ± 1.8 ± 1.2**      177      LEES      18E BABR    10.6 e<sup>+</sup>e<sup>-</sup> → π<sup>+</sup>π<sup>-</sup>3π<sup>0</sup> γ

**Γ(π<sup>+</sup>π<sup>-</sup>4π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>25</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**3.3 ± 2.3 ± 0.5**      18      LEES      21C BABR    e<sup>+</sup>e<sup>-</sup> → γ<sub>ISR</sub>(π<sup>+</sup>π<sup>-</sup>4π<sup>0</sup>)

**Γ(ρ<sup>±</sup>π<sup>∓</sup>π<sup>0</sup>π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>26</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      CL%      DOCUMENT ID      TECN      COMMENT

**<6.2**      90      LEES      18E BABR    10.6 e<sup>+</sup>e<sup>-</sup> → π<sup>+</sup>π<sup>-</sup>3π<sup>0</sup> γ

**Γ(2(π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>)) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>28</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**11.2 ± 3.3 ± 1.3**      43      AUBERT      06D BABR    10.6 e<sup>+</sup>e<sup>-</sup> → 2(π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) γ

**Γ(2(π<sup>+</sup>π<sup>-</sup>)3π<sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>30</sub>Γ<sub>7</sub>/Γ**

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

**33 ± 5 ± 5**      14k      LEES      21      BABR    10.6 e<sup>+</sup>e<sup>-</sup> → 2(π<sup>+</sup>π<sup>-</sup>)3π<sup>0</sup> γ

$\Gamma(\eta 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{33} \Gamma_7 / \Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.87 ± 1.41 ± 0.01</b>		16	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	14k	<sup>2</sup> LEES	21 BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$
----	----	-----	-------------------	---------	---

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> LEES 21 reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] < 2.3$  eV which we divide by our best value  $B(\eta \rightarrow 3\pi^0) = 32.57 \times 10^{-2}$ .

$\Gamma(\eta \pi^+ \pi^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{34} \Gamma_7 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.85</b>	90	LEES	18E BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$

$\Gamma(\eta \pi^+ \pi^- 3\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{35} \Gamma_7 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	<sup>1</sup> LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 3\pi^0 \gamma \gamma)$

<sup>1</sup> LEES 21C reports  $[\Gamma(\psi(2S) \rightarrow \eta \pi^+ \pi^- 3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

$\Gamma(\eta 2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{36} \Gamma_7 / \Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	14k	<sup>1</sup> LEES	21 BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$

<sup>1</sup> LEES 21 reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^- \pi^0)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{41} \Gamma_7 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.01 ± 0.84 ± 0.02</b>	37	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 2.69 \pm 0.73 \pm 0.16$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{42} \Gamma_7 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.2 ± 5.6 ± 0.1</b>	14k	<sup>1</sup> LEES	21 BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$

<sup>1</sup> LEES 21 reports  $[\Gamma(\psi(2S) \rightarrow \omega \pi^+ \pi^- 2\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 18 \pm 4 \pm 3$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{45}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.58±0.82±0.02</b>	33	<sup>1</sup> LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

<sup>1</sup> LEES 18E reports [ $\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ )] = 2.3 ± 0.7 ± 0.2 eV which we divide by our best value B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ ) = (89.2 ± 0.7) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega 3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{46}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.8</b>	90	<sup>1</sup> LEES	21C BABR	e <sup>+</sup> e <sup>-</sup> → γ <sub>ISR</sub> (π <sup>+</sup> π <sup>-</sup> 4π <sup>0</sup> )

<sup>1</sup> LEES 21C reports [ $\Gamma(\psi(2S) \rightarrow \omega 3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ )] < 1.6 eV which we divide by our best value B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ ) = 89.2 × 10<sup>-2</sup>.

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{51}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.55±0.19±0.01</b>	19	<sup>1</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.57±0.23±0.01	10	<sup>2</sup> AUBERT, BE	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
----------------	----	-------------------------	----------	--

<sup>1</sup> LEES 12F reports [ $\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\phi(1020) \rightarrow K^+K^-$ )] = 0.27 ± 0.09 ± 0.02 eV which we divide by our best value B( $\phi(1020) \rightarrow K^+K^-$ ) = (49.1 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Superseded by LEES 12F. AUBERT, BE 06D reports [ $\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\phi(1020) \rightarrow K^+K^-$ )] = 0.28 ± 0.11 ± 0.02 eV which we divide by our best value B( $\phi(1020) \rightarrow K^+K^-$ ) = (49.1 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{52}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.346±0.129±0.004</b>	12	<sup>1</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.346±0.168±0.004	6 ± 3	<sup>2</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
-------------------	-------	---------------------	-----------	--

<sup>1</sup> LEES 12F reports [ $\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\phi(1020) \rightarrow K^+K^-$ )] = 0.17 ± 0.06 ± 0.02 eV which we divide by our best value B( $\phi(1020) \rightarrow K^+K^-$ ) = (49.1 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports [ $\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\phi(1020) \rightarrow K^+K^-$ )] = 0.17 ± 0.08 ± 0.02 eV which we divide by our best value B( $\phi(1020) \rightarrow K^+K^-$ ) = (49.1 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{59}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.147 \pm 0.035 \pm 0.005$	66	<sup>1</sup> LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$0.197 \pm 0.035 \pm 0.005$	66	<sup>2</sup> LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$0.35 \pm 0.14 \pm 0.03$	11	<sup>3</sup> LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$

<sup>1</sup>  $\sin\phi > 0$ .<sup>2</sup>  $\sin\phi < 0$ .<sup>3</sup> Interference with non-resonant  $K^+K^-$  production not taken into account. $\Gamma(K^+K^-\pi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{60}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>1.92 \pm 0.30 \pm 0.06</math></b>	133	LEES 12F	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
--	-----	----------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.56 \pm 0.42 \pm 0.16$	85	<sup>1</sup> AUBERT 07AK	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
--------------------------	----	--------------------------	------	--

<sup>1</sup> Superseded by LEES 12F. $\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{64}\Gamma_7/\Gamma$ 

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

<b>&lt;0.7</b>	90	8	LEES 17A	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$
----------------	----	---	----------	------	---

 $\Gamma(K^+K^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{65}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>0.60 \pm 0.31 \pm 0.03</math></b>	17	LEES 12F	BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0 K^+K^-\gamma$
--	----	----------	------	---

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{66}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>4.4 \pm 1.3 \pm 0.3</math></b>	32	AUBERT 07AU	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$
---	----	-------------	------	---

 $\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{73}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>2.92 \pm 1.27 \pm 0.15</math></b>	14	LEES 17A	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$
--	----	----------	------	---

 $\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{74}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>3.14 \pm 1.08 \pm 0.16</math></b>	16	LEES 17A	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \eta \gamma$
--	----	----------	------	--

 $\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{77}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b><math>3.05 \pm 1.80 \pm 0.01</math></b>	7	<sup>1</sup> AUBERT 07AU	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$
--	---	--------------------------	------	--

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{78} \Gamma_{\gamma} / \Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.4 ± 2.1 ± 0.3</b>	26	AUBERT	06D	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

 $\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{81} \Gamma_{\gamma} / \Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.22 ± 0.10 ± 0.02</b>	13	LEES	12F	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

 $\Gamma(p \bar{p}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{101} \Gamma_{\gamma} / \Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.686 ± 0.019 OUR FIT</b>				

**0.63 ± 0.05 OUR AVERAGE** Error includes scale factor of 1.2.

0.67 ± 0.12 ± 0.02	43	<sup>1</sup> LEES	130	BABR $e^+ e^- \rightarrow p \bar{p} \gamma$
0.74 ± 0.07 ± 0.04	142	<sup>2</sup> LEES	13Y	BABR $e^+ e^- \rightarrow p \bar{p} \gamma$
0.579 ± 0.038 ± 0.036	2.7k	ANDREOTTI	07 E835	$p \bar{p} \rightarrow e^+ e^-, J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.70 ± 0.17 ± 0.03	22	<sup>3</sup> AUBERT	06B	BABR $e^+ e^- \rightarrow p \bar{p} \gamma$

<sup>1</sup> ISR photon reconstructed in the detector<sup>2</sup> ISR photon undetected<sup>3</sup> Superseded by LEES 130 $\Gamma(\Lambda \bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{124} \Gamma_{\gamma} / \Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>1.5 ± 0.4 ± 0.1</b>	AUBERT	07BD	BABR 10.6 $e^+ e^- \rightarrow \Lambda \bar{\Lambda} \gamma$

 $\psi(2S)$  BRANCHING RATIOS $\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$   $\Gamma_1 / \Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.9785 ± 0.0013 OUR AVERAGE</b>			
0.9779 ± 0.0015	<sup>1</sup> BAI	02B	BES2 $e^+ e^-$
0.981 ± 0.003	<sup>1</sup> LUTH	75	MRK1 $e^+ e^-$

<sup>1</sup> Includes cascade decay into  $J/\psi(1S)$ . $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$   $\Gamma_2 / \Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0173 ± 0.0014 OUR AVERAGE</b>			Error includes scale factor of 1.5.
0.0166 ± 0.0010	<sup>1,2</sup> SETH	04	RVUE $e^+ e^-$
0.0199 ± 0.0019	<sup>1</sup> BAI	02B	BES2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.029 ± 0.004	<sup>1</sup> LUTH	75	MRK1 $e^+ e^-$
---------------	-------------------	----	----------------

<sup>1</sup> Included in  $\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ .<sup>2</sup> Using  $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$  from RPP-2002 and  $R = 2.28 \pm 0.04$  determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(g g g)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>10.58 ± 1.62</b>	2.9 M	<sup>1</sup> LIBBY	09 CLEO	$\psi(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using  $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$  from LIBBY 09,  $B(\psi(2S) \rightarrow X J/\psi)$  relative and absolute branching fractions from MENDEZ 08,  $B(\psi(2S) \rightarrow \gamma \eta_c)$  from MITCHELL 09, and  $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$ ,  $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$ , and  $B(\psi(2S) \rightarrow \ell^+ \ell^-)$  from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  LIBBY 09 measurement.

 $\Gamma(\gamma g g)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.025 ± 0.288</b>	200 k	<sup>1</sup> LIBBY	09 CLEO	$\psi(2S) \rightarrow \gamma + \text{hadrons}$

<sup>1</sup> Calculated using  $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$  from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of  $\Gamma(g g g)/\Gamma_{\text{total}}$  LIBBY 09 measurement.

 $\Gamma(\gamma g g)/\Gamma(g g g)$   $\Gamma_4/\Gamma_3$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.7 ± 2.6 ± 1.6</b>	2.9 M	LIBBY	09 CLEO	$\psi(2S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.154 ± 0.015</b>	<sup>1</sup> MENDEZ	08 CLEO	$e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.169 ± 0.026	<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S)$
---------------	-------------------	----------	--------------------------------

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow J/\psi X)$  from MENDEZ 08 and other branching fractions from PDG 07.

<sup>2</sup> Uses  $B(J/\psi X)$  from ADAM 05A,  $B(\chi_{cJ} \gamma)$ ,  $B(\eta_c \gamma)$  from ATHAR 04 and  $B(\ell^+ \ell^-)$  from PDG 04. Superseded by MENDEZ 08.

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>79.3 ± 1.7 OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13	<sup>1</sup> FELDMAN	77 RVUE	$e^+ e^-$
---------	----------------------	---------	-----------

<sup>1</sup> From an overall fit assuming equal partial widths for  $e^+ e^-$  and  $\mu^+ \mu^-$ . For a measurement of the ratio see the entry  $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$  below. Includes LUTH 75, HILGER 75, BURMESTER 77.

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>
<b>80 ± 6 OUR FIT</b>	

 $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$   $\Gamma_8/\Gamma_7$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.00 ± 0.08 OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16	BOYARSKI	75C MRK1	$e^+ e^-$
-------------	----------	----------	-----------



$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$				$\Gamma_9/\Gamma$
VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT	
<b>31 ± 4 OUR FIT</b>				
<b>30.8 ± 2.1 ± 3.8</b>	<sup>1</sup> ABLIKIM	06W BES	$e^+e^- \rightarrow \psi(2S)$	

<sup>1</sup> Computed using PDG 02 value of  $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$  to estimate the total number of  $\psi(2S)$  events.

———— DECAYS INTO  $J/\psi(1S)$  AND ANYTHING ————

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$				
$\Gamma_{10}/\Gamma = (\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{163} + 0.190\Gamma_{164})/\Gamma$				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.614 ± 0.006 OUR FIT</b>				
<b>0.55 ± 0.07 OUR AVERAGE</b>				
0.51 ± 0.12		BRANDELIK	79C DASP	$e^+e^- \rightarrow \mu^+\mu^-X$
0.57 ± 0.08		ABRAMS	75B MRK1	$e^+e^- \rightarrow \mu^+\mu^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.644 ± 0.006 ± 0.016		<sup>1</sup> ABLIKIM	21Z BES3	$e^+e^- \rightarrow \ell^+\ell^-X$
0.6254 ± 0.0016 ± 0.0155	1.1M	<sup>2</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \ell^+\ell^-X$
0.5950 ± 0.0015 ± 0.0190	151k	ADAM	05A CLEO	Repl. by MENDEZ 08


<sup>1</sup> From a fit to the  $e^+e^- \rightarrow J/\psi X$  cross section between 3.645 and 3.891 GeV, with  $\Gamma(ee)$  and  $\Gamma$  fixed to the PDG 20 values of the cross particle fit which are correlated to "OUR FIT" value for  $B(\psi(2S) \rightarrow J/\psi X)$ .

<sup>2</sup> Not independent from other measurements of MENDEZ 08.

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$				$\Gamma_7/\Gamma_{10}$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.291 ± 0.026 OUR FIT</b>				
<b>1.28 ± 0.04 OUR AVERAGE</b>	Error includes scale factor of 1.6. See the ideogram below.			
1.22 ± 0.02 ± 0.05	5097 ± 73	<sup>1</sup> ANDREOTTI	05 E835	$\rho\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$
1.28 ± 0.03 ± 0.02		<sup>1</sup> AMBROGIANI	00A E835	$\rho\bar{p} \rightarrow \psi(2S)$
1.44 ± 0.08 ± 0.02		<sup>1</sup> ARMSTRONG	97 E760	$\bar{p}p \rightarrow \psi(2S)$

<sup>1</sup> Using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

WEIGHTED AVERAGE  
 $1.28 \pm 0.04$  (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$   $\Gamma_7/\Gamma_{10}$   
 (units  $10^{-2}$ )

**$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\text{anything})$**   **$\Gamma_8/\Gamma_{10}$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.0130 \pm 0.0010</math> OUR FIT</b>			
<b><math>0.014 \pm 0.003</math></b>	HILGER	75	SPEC $e^+e^-$

**$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$**   **$\Gamma_{11}/\Gamma$**

VALUE	DOCUMENT ID
<b><math>0.2538 \pm 0.0032</math> OUR FIT</b>	

**$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{12}/\Gamma$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------


**$0.3468 \pm 0.0030$  OUR FIT**  
 **$0.348 \pm 0.005$  OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

$0.3498 \pm 0.0002 \pm 0.0045$	20M	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$0.3504 \pm 0.0007 \pm 0.0077$	565k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
$0.323 \pm 0.014$		BAI	02B	BES2	$e^+e^-$
$0.32 \pm 0.04$		ABRAMS	75B	MRK1	$e^+e^- \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.3354 \pm 0.0014 \pm 0.0110$	60k	<sup>1</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
--------------------------------	-----	-------------------	-----	------	--------------------

WEIGHTED AVERAGE  
 $0.348 \pm 0.005$  (Error scaled by 1.3)



<sup>1</sup> Not independent from other values reported by ADAM 05A.  
 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_7/\Gamma_{12}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0229 ± 0.0005 OUR FIT</b>			
<b>0.0252 ± 0.0028 ± 0.0011</b>	<sup>1</sup> AUBERT	02B	BABR $e^+e^-$

<sup>1</sup> Using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_8/\Gamma_{12}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0230 ± 0.0017 OUR FIT</b>			
<b>0.0228 ± 0.0018 OUR AVERAGE</b>			
0.0230 ± 0.0020 ± 0.0012	<sup>1</sup> AAIJ	16Y	LHCB $\Lambda_b^0 \rightarrow \psi(2S)X$
0.0216 ± 0.0026 ± 0.0014	<sup>2</sup> AUBERT	02B	BABR $e^+e^-$
0.0327 ± 0.0077 ± 0.0072	<sup>2</sup> GRIBUSHIN	96	FMPS 515 $\pi^- \text{Be} \rightarrow 2\mu X$

<sup>1</sup> Using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$ .

<sup>2</sup> Using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$ .

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_9/\Gamma_{12}$


VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.8 ± 1.1 OUR FIT</b>			
<b>8.73 ± 1.39 ± 1.57</b>	BAI	02	BES $e^+e^-$

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma_{12}/\Gamma_{10}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.5645 ± 0.0026 OUR FIT</b>				
<b>0.554 ± 0.008 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
0.5604 ± 0.0009 ± 0.0062	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	<sup>1,2</sup> ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97	E760 $\bar{p}p \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.5637 ± 0.0027 ± 0.0046	60k	ADAM	05A	CLEO Repl. by MENDEZ 08

WEIGHTED AVERAGE  
0.554 ± 0.008 (Error scaled by 1.3)



<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> ABLIKIM 04B quotes  $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$ .

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{11}/\Gamma_{12} = (0.9761\Gamma_{13} + 0.719\Gamma_{14} + 0.343\Gamma_{163} + 0.190\Gamma_{164})/\Gamma_{12}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.732 ± 0.008 OUR FIT</b>			
<b>0.73 ± 0.09</b>	TANENBAUM 76	MRK1	$e^+ e^-$

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{13}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.1824 ± 0.0031 OUR FIT</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.1769 \pm 0.0008 \pm 0.0053$	61k	<sup>1</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
$0.1652 \pm 0.0014 \pm 0.0058$	13.4k	<sup>2</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

<sup>2</sup> Not independent from other values reported by ADAM 05A.

### $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma_{13}/\Gamma_{10}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.2968 ± 0.0031 OUR FIT**

**0.320 ± 0.012 OUR AVERAGE**

$0.300 \pm 0.008 \pm 0.022$	$1655 \pm 44$	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
$0.328 \pm 0.013 \pm 0.008$		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
$0.323 \pm 0.033$		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.2829 \pm 0.0012 \pm 0.0056$	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
$0.2776 \pm 0.0025 \pm 0.0043$	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

### $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{13}/\Gamma_{12}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.526 ± 0.008 OUR FIT**

**0.513 ± 0.022 OUR AVERAGE** Error includes scale factor of 2.2.

$0.5047 \pm 0.0022 \pm 0.0102$	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
$0.570 \pm 0.009 \pm 0.026$	14k	<sup>1</sup> ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.4924 \pm 0.0047 \pm 0.0086$	73k	<sup>2,3</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
$0.571 \pm 0.018 \pm 0.044$		<sup>4</sup> ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$	
$0.53 \pm 0.06$		TANENBAUM 76	MRK1	$e^+ e^-$	
$0.64 \pm 0.15$		<sup>5</sup> HILGER 75	SPEC	$e^+ e^-$	

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> Not independent from other values reported by ADAM 05A.

<sup>3</sup> Using 13,217  $J/\psi\pi^0\pi^0$  and 60,010  $J/\psi\pi^+\pi^-$  events.

<sup>4</sup> Not independent from other values reported by ANDREOTTI 05.

<sup>5</sup> Ignoring the  $J/\psi(1S)\eta$  and  $J/\psi(1S)\gamma\gamma$  decays.

### $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$

$\Gamma_{14}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**33.7 ± 0.5 OUR FIT**


**32.9 ± 1.7 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

$33.75 \pm 0.17 \pm 0.86$	68.2k	ABLIKIM	12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
$29.8 \pm 0.9 \pm 2.3$	5.7k	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
$25.5 \pm 2.9$	386	<sup>1</sup> OREGLIA	80	CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
$45 \pm 12$	17	<sup>2</sup> BRANDELIK	79B	DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
$42 \pm 6$	164	<sup>2</sup> BARTEL	78B	CNTR	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$34.3 \pm 0.4 \pm 0.9$	18.4k	<sup>3</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
$32.5 \pm 0.6 \pm 1.1$	2.8k	<sup>4</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
$43 \pm 8$	44	TANENBAUM 76	MRK1	$e^+ e^-$	

WEIGHTED AVERAGE  
 $32.9 \pm 1.7$  (Error scaled by 2.1)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

- <sup>1</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .
- <sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .
- <sup>3</sup> Not independent from other measurements of MENDEZ 08.
- <sup>4</sup> Not independent from other values reported by ADAM 05A.


$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$  (units  $10^{-3}$ )

**$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$**

**$\Gamma_{14}/\Gamma_{10}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.0549 \pm 0.0008</math></b>				<b>OUR FIT</b>
<b><math>0.058 \pm 0.007</math></b>				<b>OUR AVERAGE</b> Error includes scale factor of 1.4. See the ideogram below.
$0.050 \pm 0.006 \pm 0.003$	$298 \pm 20$	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
$0.072 \pm 0.009$		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
$0.061 \pm 0.015$		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
$0.0549 \pm 0.0006 \pm 0.0009$	18.4k	<sup>1</sup> MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
$0.0546 \pm 0.0010 \pm 0.0007$	2.8k	ADAM 05A	CLEO	Repl. by MENDEZ 08

WEIGHTED AVERAGE  
 $0.058 \pm 0.007$  (Error scaled by 1.4)



<sup>1</sup> Not independent from other measurements of MENDEZ 08.  
 $\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

**$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_{14}/\Gamma_{12}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.0972 \pm 0.0014</math></b>				<b>OUR FIT</b>
<b><math>0.0979 \pm 0.0018</math></b>				<b>OUR AVERAGE</b>
$0.0979 \pm 0.0010 \pm 0.0015$	18.4k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\eta$
$0.098 \pm 0.005 \pm 0.010$	2k	<sup>1</sup> ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
$0.091 \pm 0.021$		<sup>2</sup> HIMEL 80	MRK2	$e^+e^- \rightarrow \psi(2S)X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.0968 \pm 0.0019 \pm 0.0013$	2.8k	<sup>3</sup> ADAM 05A	CLEO	Repl. by MENDEZ 08
$0.095 \pm 0.007 \pm 0.007$		<sup>4</sup> ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> The value for  $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$  reported in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$ .

<sup>3</sup> Not independent from other values reported by ADAM 05A.

<sup>4</sup> Not independent from other values reported by ANDREOTTI 05.

**$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>12.68 \pm 0.32</math></b>				<b>OUR AVERAGE</b>
$12.6 \pm 0.2 \pm 0.3$	4.1k	ABLIKIM 12M	BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$
$13.3 \pm 0.8 \pm 0.3$	530	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-2\gamma$
$14.3 \pm 1.4 \pm 1.2$	280	BAI 04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
$14 \pm 6$	7	HIMEL 80	MRK2	$e^+e^-$
$9 \pm 2 \pm 1$	23	<sup>1</sup> OREGLIA 80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13 ±1 ±1 88 ADAM 05A CLEO Repl. by MENDEZ 08

<sup>1</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

### $\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_{15}/\Gamma_{10} = \Gamma_{15}/(\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{163} + 0.190\Gamma_{164})$$

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.213 ± 0.012 ± 0.003	527	<sup>1</sup> MENDEZ	08 CLEO	$e^+ e^- \rightarrow J/\psi \gamma \gamma$
0.22 ± 0.02 ± 0.01		<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$

<sup>1</sup> Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

<sup>2</sup> Not independent from other values reported by ADAM 05A.

### $\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+ \pi^-)$

$\Gamma_{15}/\Gamma_{12}$

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.380 ± 0.022 ± 0.005	527	<sup>1</sup> MENDEZ	08 CLEO	$e^+ e^- \rightarrow J/\psi \gamma \gamma$
0.39 ± 0.04 ± 0.01		<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$

<sup>1</sup> Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

<sup>2</sup> Not independent from other values reported by ADAM 05A.

## ———— HADRONIC DECAYS ————

### $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

$\Gamma_{16}/\Gamma$

VALUE (units 10 <sup>-5</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

**0.78 ± 0.26 OUR AVERAGE**

0.76 ± 0.25 ± 0.06	30	<sup>1</sup> METREVELI	12		$\psi(2S) \rightarrow \pi^+ \pi^-$
8 ± 5		BRANDELIK	79C DASP		$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	DOBBS	06A CLEO		$e^+ e^- \rightarrow \psi(2S)$
<5	90	FELDMAN	77 MRK1		$e^+ e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using  $\psi(3770) \rightarrow \pi^+ \pi^-$  for continuum subtraction.

### $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

$\Gamma_{17}/\Gamma$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------


**2.01 ± 0.17 OUR AVERAGE** Error includes scale factor of 1.7. See the ideogram below.

2.14 ± 0.03 <sup>+0.12</sup> <sub>-0.11</sub>	7k	<sup>1</sup> ABLIKIM	12H BES3	$e^+ e^- \rightarrow \psi(2S)$
1.81 ± 0.18 ± 0.19	260 ± 19	<sup>2</sup> ABLIKIM	05J BES2	$e^+ e^- \rightarrow \psi(2S)$
1.88 <sup>+0.16</sup> <sub>-0.15</sub> ± 0.28	194	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
0.85 ± 0.46	4	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

<sup>1</sup> From  $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$  events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of  $\psi(2S)$  events.

<sup>2</sup> From a PW analysis of  $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$ .





**$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{18}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.32 \pm 0.12</math></b>					<b>OUR AVERAGE</b> Error includes scale factor of 1.8.
$0.51 \pm 0.07 \pm 0.11$			<sup>1</sup> ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
$0.24^{+0.08}_{-0.07} \pm 0.02$		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83 MRK2	$e^+e^-$
<10	90		BARTEL	76 CNTR	$e^+e^-$
<10	90		<sup>2</sup> ABRAMS	75 MRK1	$e^+e^-$

<sup>1</sup> From a PW analysis of  $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ .

<sup>2</sup> Final state  $\rho^0\pi^0$ .

**$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{19}/\Gamma$**

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.94 \pm 0.25^{+1.15}_{-0.34}</math></b>	<sup>1</sup> ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From a PW analysis of  $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ .

**$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$**   **$\Gamma_{20}/\Gamma$**


VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.4 \pm 0.6</math></b>				<b>OUR AVERAGE</b> Error includes scale factor of 2.2.
$2.2 \pm 0.2 \pm 0.2$	308	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$
$4.5 \pm 1.0$		TANENBAUM	78 MRK1	$e^+e^-$

**$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{21}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.2 \pm 0.6</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.4.		
$2.0 \pm 0.2 \pm 0.4$	285.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
$4.2 \pm 1.5$		TANENBAUM	78	MRK1 $e^+ e^-$

**$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{22}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>29 \pm 10</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 4.7. See the ideogram below.		
$24.9 \pm 0.7 \pm 3.6$	2173	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$
$127 \pm 12 \pm 2$	410	<sup>1</sup> AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$
$26.1 \pm 0.7 \pm 3.0$	1703	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$
$30 \pm 8$	42	FRANKLIN	83	MRK2 $e^+ e^-$



<sup>1</sup>AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$  (units  $10^{-4}$ )

**$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$**   **$\Gamma_{23}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.55 \pm 0.73 \pm 0.47</math></b>		$112 \pm 31$	BAI	04C	BES2 $\psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.3		90	BAI	98J	BES $e^+ e^-$

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.5 ± 2.0 OUR AVERAGE</b>				Error includes scale factor of 2.8.
5.45 ± 0.42 ± 0.87	671	ABLIKIM 05H	BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+\pi^-)$
1.5 ± 1.0		<sup>1</sup> TANENBAUM 78	MRK1	$e^+e^-$

<sup>1</sup> Assuming entirely strong decay.

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>35 ± 16</b>	6	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.6</b>	90	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$

$\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.5 ± 0.7 ± 1.5</b>		<sup>1</sup> BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
10.3 ± 0.8 ± 1.4	201.7	<sup>2</sup> BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$
8.1 ± 1.4 ± 1.6	50.0	<sup>2</sup> BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$

<sup>1</sup> Average of  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow 3\pi$ .

<sup>2</sup> Not independent from other values reported by BRIERE 05.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2 ± 0.6 OUR AVERAGE</b>				Error includes scale factor of 1.1.
3.0 <sup>+1.1</sup> / <sub>-0.9</sub> ± 0.2	18	ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$
1.78 <sup>+0.67</sup> / <sub>-0.62</sub> ± 0.17	13	ABLIKIM 04L	BES	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.5 ± 1.6 ± 1.3</b>	12.8	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

$\Gamma(\eta'\rho)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.87 <sup>+1.64</sup>/<sub>-1.11</sub> ± 0.33</b>	2	ABLIKIM 04L	BES	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


1.02 ± 0.11 ± 0.24	143	<sup>1</sup> ABLIKIM 17AK	BES3	$e^+e^- \rightarrow \psi(2S)$
--------------------	-----	---------------------------	------	-------------------------------

0.569±0.128±0.236      80      <sup>2</sup> ABLIKIM      17AK BES3      e<sup>+</sup>e<sup>-</sup> → ψ(2S)  
<sup>1</sup> Destructive-interference solution of a partial wave analysis of the decay ψ(2S) → π<sup>+</sup>π<sup>-</sup>η'.  
<sup>2</sup> Constructive-interference solution of a partial wave analysis of the decay ψ(2S) → π<sup>+</sup>π<sup>-</sup>η'.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{40}/\Gamma$
VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.1 ± 0.6 OUR AVERAGE</b>					
2.5 <sup>+1.2</sup> <sub>-1.0</sub> ± 0.2	14	ADAM	05	CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
1.87 <sup>+0.68</sup> <sub>-0.62</sub> ± 0.28	14	ABLIKIM	04L	BES	e <sup>+</sup> e <sup>-</sup> → ψ(2S)

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma$
VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>7.3±1.2 OUR AVERAGE</b> Error includes scale factor of 2.1. See the ideogram below.					
8.4±0.5±1.2	386	ABLIKIM	07D	BES2	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
12.2±2.2±0.7	37	<sup>1</sup> AUBERT	07AU	BABR	10.6 e <sup>+</sup> e <sup>-</sup> → ωπ <sup>+</sup> π <sup>-</sup> γ
8.2±0.5±0.7	391	BRIERE	05	CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S) → 2(π <sup>+</sup> π <sup>-</sup> )π <sup>0</sup>
4.8±0.6±0.7	100 ± 22	<sup>2</sup> BAI	03B	BES	ψ(2S) → 2(π <sup>+</sup> π <sup>-</sup> )π <sup>0</sup>

WEIGHTED AVERAGE  
 7.3±1.2 (Error scaled by 2.1)



<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$  eV.  
<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .  
 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$  (units 10<sup>-4</sup>)

$\Gamma(b_1^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.0 ± 0.6 OUR AVERAGE</b>		Error includes scale factor of 1.1.		
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
4.18 <sup>+0.43</sup> <sub>-0.42</sub> ± 0.92	170	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	<sup>1,2</sup> BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.8 ± 1.0		<sup>1</sup> BAI	99C BES	Repl. by BAI 03B

<sup>1</sup> Assuming  $B(b_1 \rightarrow \omega \pi) = 1$ .<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ . $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2 ± 0.4 OUR AVERAGE</b>					
2.3 ± 0.5 ± 0.4		57	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
2.05 ± 0.41 ± 0.38		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.5	90		<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
<1.7	90		BAI	98J BES	Repl. by BAI 03B

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ . $\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.25<sup>+0.47</sup><sub>-0.42</sub> ± 0.40</b>	45	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\omega \eta)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.1</b>	90	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.1	90	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$


 $\Gamma(\omega \eta')/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2<sup>+2.4</sup><sub>-2.0</sub> ± 0.7</b>	4	<sup>1</sup> ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> Calculated combining  $\eta' \rightarrow \gamma \rho$  and  $\eta \pi^+ \pi^-$  channels. $\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.04</b>	90	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
<0.4	90	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{51}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.18±0.26 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.			
2.3 ±0.8 ±0.1	19 ± 6	LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
0.9 ±0.2 ±0.1	47.6	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$	
1.5 ±0.2 ±0.2	51.5 ± 8.3	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+K^-\pi^+\pi^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.45±0.96±0.04	10 ± 4	<sup>2,3</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	



<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Using  $B(\phi \rightarrow K^+K^-) = (49.3 \pm 0.6)\%$ .

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$  (units  $10^{-4}$ )

$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{52}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.75±0.33 OUR AVERAGE</b>		Error includes scale factor of 1.6.			
1.5 ±0.5 ±0.1	12 ± 4	LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
0.6 ±0.2 ±0.1	18.4 ± 6.4	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+K^-\pi^+\pi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$1.46 \pm 0.71 \pm 0.02 \quad 6 \pm 3 \quad {}^{2,3} \text{AUBERT} \quad 07\text{AK BABR} \quad 10.6 \frac{e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma}{\pi^+ \pi^- K^+ K^- \gamma}$$

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Using  $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$ .

**$\Gamma(\phi\eta)/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.10 ± 0.31 OUR AVERAGE</b>				
3.14 ± 0.23 ± 0.23	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1} \pm 0.4$	6	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

**$\Gamma(\eta\phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2.2 × 10<sup>-6</sup></b>	90	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta\phi f_0(980)$

**$\Gamma(\phi\eta')/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.54 ± 0.20 OUR AVERAGE</b>				
1.51 ± 0.16 ± 0.12	201	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
3.1 ± 1.4 ± 0.7	8	<sup>1</sup> ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> Calculated combining  $\eta' \rightarrow \gamma\rho$  and  $\eta\pi^+\pi^-$  channels.

**$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.0 ± 0.4 ± 1.3</b>	234	<sup>1</sup> ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> ABLIKIM 19BA reports  $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$  which we divide by our best value  $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta) / \Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$**

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.46 ± 1.37 ± 0.92</b>	195	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$

**$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.44 ± 0.12 ± 0.11</b>	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+ K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$< 0.45 \quad 90 \quad \text{BAI} \quad 98\text{J BES} \quad e^+ e^- \rightarrow 2(K^+ K^-)$$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>7.48 \pm 0.23 \pm 0.39</math></b>		1.3k	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$6.2 \pm 1.5 \pm 0.2$		66	<sup>2,3</sup> LEES	15J BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
$8.3 \pm 1.5 \pm 0.2$		66	<sup>3,4</sup> LEES	15J BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
$6.3 \pm 0.6 \pm 0.3$			<sup>5</sup> DOBBS	06A CLEO	$e^+ e^-$
$10 \pm 7$			<sup>5</sup> BRANDELIK	79C DASP	$e^+ e^-$
$< 5$	90		FELDMAN	77 MRK1	$e^+ e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup>  $\sin\phi > 0$ .<sup>3</sup> Using  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = (2.37 \pm 0.04)$  keV.<sup>4</sup>  $\sin\phi < 0$ .<sup>5</sup> Interference with non-resonant  $K^+ K^-$  production not taken into account. $\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>7.3 \pm 0.5</math> OUR AVERAGE</b>				
$8.1 \pm 1.3 \pm 0.3$	133	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$7.1 \pm 0.3 \pm 0.4$	817.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
$16 \pm 4$		<sup>1</sup> TANENBAUM	78 MRK1	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$11.0 \pm 1.9 \pm 0.2$	85	<sup>2</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Assuming entirely strong decay.<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.07 \pm 0.16 \pm 0.26</math></b>		0.9k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 8.9$	90	1	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.046</math></b>	<sup>1</sup> BAI	04D BES	$e^+ e^-$

<sup>1</sup> Forbidden by CP.



$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.34±0.33 OUR AVERAGE</b>				
5.28±0.25±0.34	478 ± 23	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow K_S^0 K_L^0$
5.8 ± 0.8 ± 0.4		DOBBS	06A CLEO	$e^+ e^-$
5.24±0.47±0.48	156 ± 14	<sup>2</sup> BAI	04B BES2	$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$ . $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.6±0.9 OUR AVERAGE</b>				
18.9±5.7±0.3	32	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$
11.7±1.0±1.5	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
12.7±0.5±1.0	711.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.9±2.0±0.9</b>				
	19	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6±1.3±1.8</b>				
	238	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.6±2.2±1.7</b>				
	133	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3±2.2±1.4</b>				
	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1±1.3±1.2</b>				
	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.20±0.25±0.37</b>	83 ± 9	ABLIKIM	050 BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^- \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2±0.2±0.4</b>	223.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.86±0.32±0.43</b>	93 ± 16	BAI	04C		$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2	90	BAI	98J BES	$e^+ e^-$
------	----	-----	---------	-----------

$\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.3±0.7±0.1</b>	7	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

<sup>1</sup>AUBERT 07AU quotes  $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$  eV.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.0±2.5±1.8</b>	65	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9 ± 0.4 OUR AVERAGE</b>					Error includes scale factor of 1.2.

3.18±0.30 <sup>+0.26</sup> <sub>-0.31</sub>	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
---	------	---------	----------	--------------------------------

2.9 <sup>+1.3</sup> <sub>-1.7</sub> ± 0.4	9.6 ± 4.2	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
---	-----------	---------	----------	--------------------------------

1.3 <sup>+1.0</sup> <sub>-0.7</sub> ± 0.3	7	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
---	---	------	---------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.4	90	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons
------	----	----------	---------	-------------------------------

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$   $\Gamma_{81}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.63±0.13 OUR AVERAGE</b>				

0.9 ± 0.4 ± 0.1	13	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
-----------------	----	------	----------	--

0.6 ± 0.1 ± 0.1	59.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
-----------------	------	--------	---------	---

$\Gamma(2(K^+ K^-) \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{82}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.2±0.2</b>	44.7	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$

$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$   $\Gamma_{83}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.70±0.16 OUR AVERAGE</b>				
0.8 ±0.2 ±0.1	36.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

0.6 ±0.2 ±0.1    16.1 ± 5.0    <sup>1</sup> BAI    03B BES     $\psi(2S) \rightarrow 2(K^+ K^-)$

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{84}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>10.0±1.8±2.1</b>	<sup>1</sup> BAI	99C BES	$e^+ e^-$

<sup>1</sup> Assuming  $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.7±2.5</b>	TANENBAUM	78 MRK1	$e^+ e^-$

 $\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.49±0.09±0.15</b>	1.8k		<sup>1</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.08±0.29±0.25	0.3k	<sup>1,2</sup> ABLIKIM	12L BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$
<13	90	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>1</sup> Excluding  $\eta\phi$ .

<sup>2</sup> Superseded by ABLIKIM 20F.

 $\Gamma(X(1750)\eta \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.8±1.0±2.6</b>	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.1</b>	90	<sup>1</sup> BAI	99C BES	$e^+ e^-$

<sup>1</sup> Assuming  $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.12±0.62<sup>+1.13</sup><sub>-0.61</sub></b>	251 ± 22	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.9±2.0 OUR AVERAGE</b>				
13.3 <sup>+2.4</sup> <sub>-2.8</sub> ±1.7	65.6 ± 9.0	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
9.2 <sup>+2.7</sup> <sub>-2.2</sub> ±0.9	25	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$   $\Gamma_{80}/\Gamma_{90}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.16 ± 0.06 OUR AVERAGE</b>			
0.22 <sup>+0.10</sup> <sub>-0.14</sub>	ABLIKIM	05I	BES2 $e^+ e^- \rightarrow \psi(2S)$
0.14 <sup>+0.08</sup> <sub>-0.06</sub>	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.62 ± 0.11 OUR AVERAGE</b>				Error includes scale factor of 1.1.
1.56 ± 0.04 ± 0.11	2.8k	ABLIKIM	14G	BES3 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
2.38 ± 0.37 ± 0.29	78	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.9 ± 0.3 ± 0.3	76.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow$ $K^+ K^- \pi^+ \pi^- \pi^0$
1.5 ± 0.3 ± 0.2	23	<sup>1</sup> BAI	03B	BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

$\Gamma(\omega K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.04 ± 0.39 ± 0.36</b>	1.5k	ABLIKIM	21AL	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0 K_S^0 K_S^0$

$\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.7 ± 2.6 OUR AVERAGE</b>				
18.9 ± 2.9 ± 2.2	396	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
22.6 ± 3.0 ± 2.4	535	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.1 ± 1.2 OUR AVERAGE</b>				
6.39 ± 1.50 ± 0.78	128	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
5.86 ± 1.61 ± 0.83	143	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>16.8 ± 2.5 ± 1.6</b>	356	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.82 ± 2.08 ± 0.72</b>	116	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.60 ± 0.27 ± 0.24</b>	109	<sup>1</sup> ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

<sup>1</sup> X(1440) compatible with  $\eta(1405)$  and  $\eta(1475)$ . A  $f_1(1420)$  is also possible.

$\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{98}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.09±0.20±0.16</b>	82	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

<sup>1</sup>  $X(1440)$  compatible with  $\eta(1405)$  and  $\eta(1475)$ . A  $f_1(1420)$  is also possible.

 $\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.302±0.098±0.027</b>	22	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

<sup>1</sup> Statistical significance  $4.5 \sigma$ . This measurement is equivalent to a limit of  $< 0.478 \times 10^{-5}$  at 90% C.L.

 $\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.125±0.070±0.013</b>	10	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

<sup>1</sup> Statistical significance  $3.2 \sigma$ . This measurement is equivalent to a limit of  $< 0.221 \times 10^{-5}$  at 90% C.L.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{101}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.94±0.08 OUR FIT</b>				
<b>3.02±0.08 OUR AVERAGE</b>				
3.05±0.02±0.12	19k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.08±0.05±0.18	4.5k	<sup>1</sup> DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.36±0.09±0.25	1.6k	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.87±0.12±0.15	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
1.4 ±0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.3 ±0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_{101}/\Gamma_{12}$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.49±0.23 OUR FIT</b>			
<b>6.98±0.49±0.97</b>	BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$   $\Gamma_{102}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.06±0.06±0.14</b>	6k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{103}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.53±0.07 OUR AVERAGE</b>				
1.65±0.03±0.15	4.5k	ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
1.54±0.06±0.06	948	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$
1.32±0.10±0.15	256	<sup>1</sup> ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 ±0.5	9	FRANKLIN	83 MRK2	$e^+ e^-$

<sup>1</sup> Computed using  $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$ .

$\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{104}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.42 \pm 0.20^{+1.78}_{-1.28}$	1.9k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{105}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3 <math>^{+1.7}_{-1.5}</math> OUR AVERAGE</b>				Error includes scale factor of 2.5.

$3.58 \pm 0.25^{+1.59}_{-0.84}$	1.1k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
---------------------------------	------	----------------------	----------	--------------------------------------

$8.1 \pm 0.7 \pm 0.3$	474	<sup>2</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$
-----------------------	-----	------------------------	---------	---------------------------------------

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

<sup>2</sup> From a fit of the  $p\bar{p}$  and  $p\pi^0$  mass distributions to a combination of  $N(1440)\bar{p}$ , a broad  $p\bar{p}$  enhancement around 2100 MeV, and two other broad, unestablished resonances.

 $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{106}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.64 \pm 0.05^{+0.22}_{-0.17}$	0.2k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.47 \pm 0.28^{+0.99}_{-0.97}$	0.7k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.76 \pm 0.28^{+1.37}_{-1.66}$	1.1k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{109}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.79 \pm 0.10^{+0.24}_{-0.71}$	0.5k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{110}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.62 \pm 0.28^{+1.12}_{-0.64}$	0.9k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

$\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{111}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.13 \pm 0.08^{+0.40}_{-0.30}$	0.8k	<sup>1</sup> ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{112}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>6.0 \pm 0.4</math> OUR AVERAGE</b>				
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$
$8 \pm 2$		<sup>1</sup> TANENBAUM	78	MRK1 $e^+e^-$

<sup>1</sup> Assuming entirely strong decay.

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{113}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.7 \pm 0.6 \pm 0.4</math></b>	30.1	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>6.0 \pm 0.4</math> OUR AVERAGE</b>				
$6.4 \pm 0.2 \pm 0.6$	679	<sup>1</sup> ABLIKIM	13S	BES3 $\psi(2S) \rightarrow \eta p\bar{p}$
$5.6 \pm 0.6 \pm 0.3$	154	<sup>1</sup> ALEXANDER	10	CLEO $\psi(2S) \rightarrow \eta p\bar{p}$
$5.8 \pm 1.1 \pm 0.7$	$44.8 \pm 8.5$	<sup>2</sup> ABLIKIM	05E	BES2 $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
$8 \pm 3 \pm 3$	9.8	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

<sup>1</sup> With  $N(1535)$  decaying to  $p\eta$ .

<sup>2</sup> Computed using  $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$ .

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}$   $\Gamma_{115}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.5^{+0.7}_{-0.6}</math> OUR AVERAGE</b>				
$5.2 \pm 0.3^{+3.2}_{-1.2}$	527	<sup>1</sup> ABLIKIM	13S	BES3 $\psi(2S) \rightarrow \eta p\bar{p}$
$4.4 \pm 0.6 \pm 0.3$	123	<sup>2</sup> ALEXANDER	10	CLEO $\psi(2S) \rightarrow \eta p\bar{p}$

<sup>1</sup> With  $N(1535)$  decaying to  $p\eta$ .

<sup>2</sup> From a fit of the  $p\bar{p}$  and  $p\eta$  distributions to a combination of  $N^*(1535)\bar{p}$  and a broad  $p\bar{p}$  enhancement around 2100 MeV.

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>7.3 \pm 0.4 \pm 0.6</math></b>	434.9	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

$\Gamma(p\bar{p}\rho^0)/\Gamma_{\text{total}}$					$\Gamma_{117}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>0.5 \pm 0.1 \pm 0.2</math></b>	61.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$	

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$					$\Gamma_{118}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>0.69 \pm 0.21</math> OUR AVERAGE</b>					
$0.6 \pm 0.2 \pm 0.2$	21.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$	
$0.8 \pm 0.3 \pm 0.1$	$14.9 \pm 0.1$	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$	

<sup>1</sup>Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

$\Gamma(p\bar{p}\eta')/\Gamma_{\text{total}}$					$\Gamma_{119}/\Gamma$
VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.10 \pm 0.10 \pm 0.08</math></b>	491	<sup>1</sup> ABLIKIM	19N BES3	$\psi(2S) \rightarrow \eta' p\bar{p}$	

<sup>1</sup>From the combination of  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$  and  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$  channels.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$					$\Gamma_{120}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>6.06 \pm 0.38 \pm 0.48</math></b>		753	ABLIKIM	19AO BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<24	90		BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
<26	90		<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+K^- p\bar{p}$

<sup>1</sup>Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

$\Gamma(\phi X(1835) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}$					$\Gamma_{121}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;1.82 \times 10^{-7}</math></b>	90	ABLIKIM	19AO BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$	

$\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$					$\Gamma_{122}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.48 \pm 0.17</math> OUR AVERAGE</b>					
$2.45 \pm 0.11 \pm 0.21$	851	ABLIKIM	06I BES2	$e^+e^- \rightarrow p\pi^- X$	
$2.52 \pm 0.12 \pm 0.22$	849	ABLIKIM	06I BES2	$e^+e^- \rightarrow \bar{p}\pi^+ X$	


$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{123}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>3.18 \pm 0.50 \pm 0.50</math></b>	$135 \pm 21$	ABLIKIM	06I BES2	$e^+e^- \rightarrow p\pi^-\pi^0 X$	

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					$\Gamma_{124}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.81 \pm 0.13</math> OUR AVERAGE</b>					Error includes scale factor of 1.4. See the ideogram below.
$3.97 \pm 0.02 \pm 0.12$		31k	ABLIKIM	17L BES3	$e^+e^- \rightarrow \Lambda\bar{\Lambda}$



$3.71 \pm 0.05 \pm 0.15$	6.5k	<sup>1</sup> DOBBS	17		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
$3.39 \pm 0.20 \pm 0.32$	337	ABLIKIM	07C	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$6.4 \pm 1.8 \pm 0.1$		<sup>2</sup> AUBERT	07BD	BABR	$10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
$3.28 \pm 0.23 \pm 0.25$	208	PEDLAR	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$3.75 \pm 0.09 \pm 0.23$	1.9k	<sup>1,3</sup> DOBBS	14		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
$1.81 \pm 0.20 \pm 0.27$	80	<sup>4</sup> BAI	01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 4	90	FELDMAN	77	MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

WEIGHTED AVERAGE  
 $3.81 \pm 0.13$  (Error scaled by 1.4)



<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> AUBERT 07BD reports  $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by DOBBS 17.

<sup>4</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  (units  $10^{-4}$ )

**$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$**

**$\Gamma_{125}/\Gamma$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 0.29	90	<sup>1</sup> ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<12	90	<sup>2</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
-----	----	----------------------	----------	-------------------------------

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.4\%$ .

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$   $\Gamma_{126}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.48±0.34±0.19</b>		60	<sup>1</sup> ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.9	90	<sup>2</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
------	----	----------------------	----------	-------------------------------

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.31\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ .

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{127}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.8±0.4±0.5</b>	73.4	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$   $\Gamma_{128}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.0±0.1±0.1</b>	74.0	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$

$\Gamma(\Lambda\bar{p}K^*(892)^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{129}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.3±0.5±0.5</b>	1011	ABLIKIM	19AU BES3	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.8±0.3±0.3</b>	45.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{131}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.81±0.11±0.14</b>	50	<sup>1</sup> ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

<sup>1</sup> Using  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$ .

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$   $\Gamma_{132}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.8±1.0±3.4</b>	157	<sup>1</sup> BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{133}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.40±0.03±0.13</b>	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

$\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{134}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.54±0.04±0.13</b>	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

$\Gamma(\Lambda\bar{\Sigma}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{135}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.60±0.31±0.59</b>	60	ABLIKIM 21L	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

 $\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{136}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.23±0.23±0.08 30 <sup>1</sup> DOBBS 17  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{137}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.67±0.13±0.12** 276 <sup>1</sup> ABLIKIM 13D BES3  $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

<sup>1</sup> Using  $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ , and  $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$ .

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$   $\Gamma_{138}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.43±0.10 OUR AVERAGE** Error includes scale factor of 1.4.

2.52±0.04±0.09 5.4k ABLIKIM 21AT BES3  $\psi(2S) \rightarrow p\pi^0\bar{p}\pi^0$

2.31±0.06±0.10 1.9k <sup>1</sup> DOBBS 17  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

2.57±0.44±0.68 35 PEDLAR 05 CLEO  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.51±0.15±0.16 281 <sup>1,2</sup> DOBBS 14  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Superseded by DOBBS 17.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{139}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.35±0.09 OUR AVERAGE** Error includes scale factor of 1.1.

2.44±0.03±0.11 7k ABLIKIM 17L BES3  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

2.22±0.05±0.11 2.6k <sup>1</sup> DOBBS 17  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

2.35±0.36±0.32 59 ABLIKIM 07C BES  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

2.63±0.35±0.21 58 PEDLAR 05 CLEO  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25±0.11±0.16 439 <sup>1,2</sup> DOBBS 14  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

1.2±0.4±0.4 8 <sup>3</sup> BAI 01 BES  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Superseded by DOBBS 17.

<sup>3</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

 $\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$   $\Gamma_{140}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.5±0.7 OUR AVERAGE**

8.4±0.5±0.5 1.5k ABLIKIM 16L BES3  $\psi(2S) \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$

11±3±3 14 <sup>1</sup> BAI 01 BES  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$   $\Gamma_{141}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.85 ± 0.6 ± 0.6</b>	1.4k	ABLIKIM	16L	BES3 $\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$   $\Gamma_{142}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.69 ± 0.05 ± 0.05</b>	2.2k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$   $\Gamma_{143}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.87 ± 0.11 OUR AVERAGE</b>			Error includes scale factor of 1.1.		
3.03 ± 0.05 ± 0.14	3.6k	<sup>1</sup> DOBBS	17		$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
2.78 ± 0.05 ± 0.14	5k	ABLIKIM	16L	BES3	$\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$
3.03 ± 0.40 ± 0.32	67	ABLIKIM	07C	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
2.38 ± 0.30 ± 0.21	63	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.66 ± 0.12 ± 0.20	548	<sup>1,2</sup> DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
0.94 ± 0.27 ± 0.15	12	<sup>3</sup> BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
< 2	90	FELDMAN	77	MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.  
<sup>2</sup> Superseded by DOBBS 17.  
<sup>3</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$   $\Gamma_{144}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.4 OUR AVERAGE</b>		Error includes scale factor of 4.2.		
2.73 ± 0.03 ± 0.13	11k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
1.97 ± 0.06 ± 0.11	1.2k	<sup>1</sup> DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
2.75 ± 0.64 ± 0.61	19	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.02 ± 0.19 ± 0.15	112	<sup>1,2</sup> DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.  
<sup>2</sup> Superseded by DOBBS 17.

$\Gamma(\Xi(1530)^0 \bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$   $\Gamma_{145}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.77 ± 0.14 ± 0.39</b>		2951	ABLIKIM	21A0	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 32	90	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
< 8.1	90	<sup>1</sup> BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\Lambda \bar{\Xi}^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{146}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.86 ± 0.27 ± 0.32</b>	236	ABLIKIM	15i	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow$ $K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

$\Gamma(\Xi(1530)^-\Xi(1530)^+)/\Gamma_{\text{total}}$   $\Gamma_{147}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.45±0.40±0.59</b>	5k	ABLIKIM	19AT BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi(1530)^-\Xi^+)/\Gamma_{\text{total}}$   $\Gamma_{148}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.0±1.1±0.4</b>	399	ABLIKIM	19AT BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi(1530)^0\Xi^0)/\Gamma_{\text{total}}$   $\Gamma_{149}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.53±0.04±0.03</b>	278	ABLIKIM	21AO BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi(1690)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{150}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.21±1.48±0.57</b>	74	ABLIKIM	15i BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $K^-\Lambda\Xi^+ + \text{c.c.}$

$\Gamma(\Xi(1820)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{151}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.03±2.94±1.22</b>	136	ABLIKIM	15i BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $K^-\Lambda\Xi^+ + \text{c.c.}$

$\Gamma(\Sigma^0\Xi^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{152}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.67±0.33±0.28</b>	142	ABLIKIM	15i BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $K^-\Sigma^0\Xi^+ + \text{c.c.}$

$\Gamma(\Omega^-\bar{\Omega}^+)/\Gamma_{\text{total}}$   $\Gamma_{153}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.66±0.30 OUR AVERAGE</b>			Error includes scale factor of 1.3.		
5.85±0.12±0.25		4k	<sup>1</sup> ABLIKIM	21E BES3	$\psi(2S) \rightarrow \Omega^-\bar{\Omega}^+ \rightarrow$ $\Lambda K^-\bar{\Lambda}K^+$
5.2 ±0.3 ±0.3		326	<sup>1,2</sup> DOBBS	17	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.7 ±0.9 ±0.5		27	<sup>1,2,3</sup> DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
<15	90		ABLIKIM	12Q BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
<16	90		PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
< 7.3	90		<sup>4</sup> BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

<sup>1</sup> Using  $B(\Omega^- \rightarrow \Lambda K^-) = (67.8 \pm 0.7)\%$  and  $B(\Lambda \rightarrow p\pi^-) = (63.9 \pm 0.5)\%$ .

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>3</sup> Superseded by DOBBS 17.

<sup>4</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

$\Gamma(\eta_c \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{154}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	PEDLAR 07	CLEO	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{155}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.6 ± 1.3 OUR AVERAGE**

9.0 ± 1.5 ± 1.3      3k      <sup>1</sup> GE      11      CLEO       $\psi(2S) \rightarrow \pi^0$  anything

8.4 ± 1.3 ± 1.0      11k      ABLIKIM      10B      BES3       $\psi(2S) \rightarrow \pi^0 h_c$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen      92<sup>+23</sup><sub>-22</sub>      ADAMS      09      CLEO       $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$

seen      1282      DOBBS      08A      CLEO       $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

seen      168 ± 40      ROSNER      05      CLEO       $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

<sup>1</sup> Assuming a width  $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$ , a measured dependence of the central value of  $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4}$ , and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{156}/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.7 × 10 <sup>-6</sup>	90	450M	ABLIKIM 18Q	BES3	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{157}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.88	90	BAI 04G	BES2	$e^+ e^-$

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$   $\Gamma_{158}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	BAI 04G	BES2	$e^+ e^-$

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$   $\Gamma_{159}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.70	90	BAI 04G	BES2	$e^+ e^-$

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$   $\Gamma_{160}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	BAI 04G	BES2	$e^+ e^-$

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$   $\Gamma_{161}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.60	90	BAI 04G	BES2	$e^+ e^-$

————— **RADIATIVE DECAYS** —————

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$   $\Gamma_{162}/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.79 ± 0.20</b>				<b>OUR FIT</b>
<b>9.33 ± 0.26</b>				<b>OUR AVERAGE</b>
9.389 ± 0.014 ± 0.332	4.7M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.22 ± 0.11 ± 0.46	72k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.9 ± 0.5 ± 0.8		<sup>1</sup> GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.2 ± 2.3		<sup>1</sup> BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$
7.5 ± 2.6		<sup>1</sup> WHITAKER	76	MRK1 $e^+e^-$

<sup>1</sup> Angular distribution  $(1+\cos^2\theta)$  assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$   $\Gamma_{163}/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.75 ± 0.24</b>				<b>OUR FIT</b>
<b>9.54 ± 0.29</b>				<b>OUR AVERAGE</b>
9.905 ± 0.011 ± 0.353	5.0M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.07 ± 0.11 ± 0.54	76k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.0 ± 0.5 ± 0.7		<sup>1</sup> GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.1 ± 1.9		<sup>2</sup> BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

<sup>1</sup> Angular distribution  $(1-0.189 \cos^2\theta)$  assumed.

<sup>2</sup> Valid for isotropic distribution of the photon.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$   $\Gamma_{162}/\Gamma_{163}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • •			We do not use the following data for averages, fits, limits, etc. • • •
1.02 ± 0.01 ± 0.07	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$

<sup>1</sup> Not independent from ATHAR 04 measurements of  $B(\gamma\chi_{cJ})$ .


$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$   $\Gamma_{164}/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.52 ± 0.20</b>				<b>OUR FIT</b>
<b>9.42 ± 0.31</b>				<b>OUR AVERAGE</b>
9.621 ± 0.013 ± 0.272	4.2M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.33 ± 0.14 ± 0.61	79k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
8.0 ± 0.5 ± 0.7		<sup>1</sup> GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.0 ± 2.0		<sup>2</sup> BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

<sup>1</sup> Angular distribution  $(1-0.052 \cos^2\theta)$  assumed.

<sup>2</sup> Valid for isotropic distribution of the photon.

WEIGHTED AVERAGE  
9.42±0.31 (Error scaled by 1.3)



$$\frac{[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}}}{(\Gamma_{162} + \Gamma_{163} + \Gamma_{164})/\Gamma_{\text{total}}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
27.6±0.3±2.0	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
	<sup>1</sup> Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$ .		

$$\frac{\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))}{\Gamma_{162}/\Gamma_{164}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.99±0.02±0.08	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
	<sup>1</sup> Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$ .		

$$\frac{\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))}{\Gamma_{164}/\Gamma_{163}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.03±0.02±0.03	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
	<sup>1</sup> Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$ .		

$$\frac{\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}}{\Gamma_{165}/\Gamma_{\text{total}}}$$

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.34 ±0.05 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
0.432±0.016±0.060		MITCHELL	09	CLEO $e^+e^- \rightarrow \gamma X$
0.32 ±0.04 ±0.06	2.5k	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$




$0.28 \pm 0.06$  <sup>2</sup> GAISER 86 CBAL  $e^+e^- \rightarrow \gamma X$

<sup>1</sup> ATHAR 04 used  $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$  MeV to obtain this result.

<sup>2</sup> GAISER 86 used  $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$  MeV to obtain this result.

WEIGHTED AVERAGE  
 $0.34 \pm 0.05$  (Error scaled by 1.3)



**$\Gamma(\gamma \eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{166}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>7 \pm 2 \pm 4</math></b>		<sup>1</sup> ABLIKIM 12G	BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 8	90	<sup>2</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
< 20	90	ATHAR 04	CLEO	$e^+e^- \rightarrow \gamma X$
20–130	95	EDWARDS 82C	CBAL	$e^+e^- \rightarrow \gamma X$

<sup>1</sup> ABLIKIM 12G reports  $[\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K \bar{K} \pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$  which we divide by our best value  $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = (1.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> CRONIN-HENNESSY 10 reports  $[\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K \bar{K} \pi)] < 14.5 \times 10^{-6}$  which we divide by our best value  $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = 1.9 \times 10^{-2}$ . This measurement assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

**$\Gamma(\gamma \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{167}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b><math>1.04 \pm 0.22</math></b>					OUR AVERAGE Error includes scale factor of 1.4.
$0.95 \pm 0.16 \pm 0.05$		423	ABLIKIM 17X	BES3	$\psi(2S) \rightarrow \gamma \pi^0$
$1.58 \pm 0.40 \pm 0.13$		37	ABLIKIM 10F	BES3	$\psi(2S) \rightarrow \gamma \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400	95	<sup>1</sup> LIBERMAN	75	SPEC	$e^+ e^-$
< $1 \times 10^4$	90	WIIK	75	DASP	$e^+ e^-$

<sup>1</sup> Restated by us using  $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$ .

**$\Gamma(\gamma 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$**   **$\Gamma_{168}/\Gamma$**

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>39.6 \pm 2.8 \pm 5.0</math></b>	583	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$

**$\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$**   **$\Gamma_{169}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 17</b>	90	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$

**$\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$**   **$\Gamma_{170}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.24 \pm 0.04</math> OUR AVERAGE</b>					
$1.251 \pm 0.022 \pm 0.062$		56k	ABLIKIM	17X	BES3 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$ , $\gamma \pi^0 \pi^0 \eta$
$1.26 \pm 0.03 \pm 0.08$		2226	<sup>1</sup> ABLIKIM	10F	BES3 $\psi(2S) \rightarrow 3\gamma \pi^+ \pi^-$ , $2\gamma \pi^+ \pi^-$
$1.19 \pm 0.08 \pm 0.03$			PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
$1.24 \pm 0.27 \pm 0.15$		23	ABLIKIM	06R	BES2 $e^+ e^- \rightarrow \psi(2S)$
$1.54 \pm 0.31 \pm 0.20$		$\sim 43$	BAI	98F	BES $\psi(2S) \rightarrow \pi^+ \pi^- 2\gamma$ , $\pi^+ \pi^- 3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60	90	<sup>2</sup> BRAUNSCH...	77	DASP	$e^+ e^-$
< 11	90	<sup>3</sup> BARTEL	76	CNTR	$e^+ e^-$

<sup>1</sup> Combining the results from  $\eta' \rightarrow \pi^+ \pi^- \eta$  and  $\eta' \rightarrow \pi^+ \pi^- \gamma$  decay modes.

<sup>2</sup> Restated by us using total decay width 228 keV.

<sup>3</sup> The value is normalized to the branching ratio for  $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ .

**$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$**   **$\Gamma_{171}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.73^{+0.29}_{-0.25}</math> OUR AVERAGE</b>				Error includes scale factor of 1.8.

$2.84 \pm 0.15^{+0.03}_{-0.10}$	1.9k	1,2	DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$
$2.12 \pm 0.19 \pm 0.32$		3,4	BAI	03C	BES $\psi(2S) \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.08 \pm 0.19 \pm 0.33$	$200.6 \pm 18.8$	<sup>3</sup>	BAI	03C	BES $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
$2.90 \pm 1.08 \pm 1.07$	$29.9 \pm 11.1$	<sup>3</sup>	BAI	03C	BES $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

<sup>4</sup> Combining the results from  $\pi^+ \pi^-$  and  $\pi^0 \pi^0$  decay modes.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{172}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1 ± 1.0 ± 1.4</b>	175	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$   $\Gamma_{173}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3 ± 1.8 ± 0.6</b>	274	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

<sup>1</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi \pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$  which we divide by our best value  $B(f_0(1500) \rightarrow \pi \pi) = (34.5 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{174}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.3 ± 0.8 ± 0.1</b>	136	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

<sup>1</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$  which we divide by our best value  $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$   $\Gamma_{176}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.5 ± 0.6 OUR AVERAGE</b>				
3.6 ± 0.4 ± 0.5	290	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$
3.01 ± 0.41 ± 1.24	35.6 ± 4.8	<sup>2</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{177}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.6 ± 0.7 OUR AVERAGE</b>					
6.7 ± 0.6 ± 0.6		375	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$
6.04 ± 0.90 ± 1.32	39.6 ± 5.9		<sup>2,3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.6	90	6.8 ± 3.1	<sup>2,3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
--------	----	-----------	--------------------	---------	---

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Includes unknown branching fractions to  $K^+ K^-$  or  $K_S^0 K_S^0$ . We have multiplied the  $K^+ K^-$  result by a factor of 2 and the  $K_S^0 K_S^0$  result by a factor of 4 to obtain the  $K \bar{K}$  result.

<sup>3</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$   $\Gamma_{178}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.8 ± 0.5 ± 0.9</b>	373	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{179}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.2 \pm 0.6 \pm 0.8</math></b>	207	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$   $\Gamma_{180}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 5.8 \times 10^{-6}</math></b>	90	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $\pi^+ \pi^-$  and  $\pi^0 \pi^0$  are  $3.2/4.3 \times 10^{-6}$  and  $2.6/4.0 \times 10^{-6}$ , respectively. $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{181}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 9.5 \times 10^{-6}</math></b>	90	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $K^+ K^-$  and  $K_S^0 K_S^0$  are  $2.1/4.3 \times 10^{-6}$  and  $3.7/5.5 \times 10^{-6}$ , respectively. $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$   $\Gamma_{182}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.92 \pm 0.18</math> OUR AVERAGE</b>					
$0.85 \pm 0.18 \pm 0.04$		382	<sup>1</sup> ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$ , $\gamma 3\pi^0$
$1.38 \pm 0.48 \pm 0.09$		13	<sup>1</sup> ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$ , $\gamma 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2$	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
$< 90$	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
$< 200$	90	YAMADA	77	DASP	$e^+ e^- \rightarrow 3\gamma$

<sup>1</sup> Combining the results from  $\eta \rightarrow \pi^+ \pi^- \pi^0$  and  $\eta \rightarrow 3\pi^0$  decay modes. $\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{183}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>8.71 \pm 1.25 \pm 1.64</math></b>	418	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$   $\Gamma_{185}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 0.9</math></b>	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.3$	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
$< 1.2$	90	<sup>1</sup> SCHARRE	80	MRK1 $e^+ e^-$

<sup>1</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow K \bar{K} \pi$ . $\Gamma(\gamma \eta(1405) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{186}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.36 \pm 0.25 \pm 0.05</math></b>	10	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{187}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0 \times 10^{-7}$	90	ABLIKIM	17AJ BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$   $\Gamma_{189}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4$	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.5$	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
--------	----	---------	----------	---

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{190}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<0.88$	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{191}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$37.0 \pm 6.1 \pm 7.2$	237	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$   $\Gamma_{192}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$24.0 \pm 4.5 \pm 5.0$	41	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{193}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$25.6 \pm 3.6 \pm 3.6$	115	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{194}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$19.1 \pm 2.7 \pm 4.3$	132	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{195}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<22$	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma 2(K^+ K^-))/\Gamma_{\text{total}}$   $\Gamma_{196}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<4$	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{197}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.9 ± 0.5 OUR AVERAGE</b>				Error includes scale factor of 2.0.
$4.18 \pm 0.26 \pm 0.18$	348	<sup>1</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
$2.9 \pm 0.4 \pm 0.4$	142	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{198}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.2 \pm 0.2 \pm 0.1</math></b>	111	<sup>1</sup> ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{199}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.72 \pm 0.18 \pm 0.03</math></b>	73	<sup>1</sup> ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{200}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>4.57 \pm 0.36^{+1.77}_{-4.26}</math></b>		ABLIKIM 12D	BES3	$J/\psi \rightarrow \gamma p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
<5.4	90	ABLIKIM 07D	BES	$\psi(2S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{201}/\Gamma$ 

For a narrow resonance in the range  $2.2 < M(X) < 2.8$  GeV.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2</b>	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma p \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{202}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.8 \pm 1.2 \pm 0.7</math></b>	17	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma \gamma J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{204}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.1 \pm 0.6^{+0.8}_{-1.0}</math></b>	1.1k	ABLIKIM 12O	BES3	$e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.2 \pm 0.6$	1.1k	<sup>1</sup> ABLIKIM 17N	BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
---------------	------	--------------------------	------	---

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ . No systematic error estimation.

 $\Gamma(e^+ e^- \eta')/\Gamma_{\text{total}}$   $\Gamma_{205}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.90 \pm 0.26</math> OUR AVERAGE</b>				
$1.99 \pm 0.33 \pm 0.12$	57	ABLIKIM 18Z	BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$ , $\eta' \rightarrow \gamma \pi^+ \pi^-$
$1.79 \pm 0.38 \pm 0.11$	20	ABLIKIM 18Z	BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$ , $\eta' \rightarrow \eta \pi^+ \pi^-$

$\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}$   $\Gamma_{206}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>10.6±2.4±0.4</b>	48	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma(\gamma \chi_{c0}(1P))$   $\Gamma_{206}/\Gamma_{162}$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.4±1.9±0.6</b>	48	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}$   $\Gamma_{207}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.5±0.6±0.2</b>	873	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma(\gamma \chi_{c1}(1P))$   $\Gamma_{207}/\Gamma_{163}$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.3±0.3±0.4</b>	873	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}$   $\Gamma_{208}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.0±0.7±0.2</b>	227	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma(\gamma \chi_{c2}(1P))$   $\Gamma_{208}/\Gamma_{164}$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.6±0.5±0.4</b>	227	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

---

**WEAK DECAYS**


---

$\Gamma(D^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{209}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-7}$	90	<sup>1</sup> ABLIKIM	17AF BES3	$e^+ e^- \rightarrow \psi(2S)$	

<sup>1</sup> Using  $D^0$  decays to  $K^- \pi^+$ ,  $K^- \pi^+ \pi^0$ , and  $K^- \pi^+ \pi^+ \pi^-$ .

---

**OTHER DECAYS**


---

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$					$\Gamma_{210}/\Gamma_7$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0$	90	LEES	13I BABR	$B \rightarrow K^{(*)} \psi(2S)$	

---

 **$\psi(2S)$  CROSS-PARTICLE BRANCHING RATIOS**

For measurements involving  $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$  see the corresponding entries in the  $\chi_{cJ}(1P)$  sections.

---

**MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS**
 **$\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)$  and  $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$** 
 **$a_2(\chi_{c1})/a_2(\chi_{c2})$  Magnetic quadrupole transition amplitude ratio**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>63 \pm 7</math> OUR AVERAGE</b>				
$61.7 \pm 8.3$	253k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$67^{+19}_{-13}$	59k	<sup>2</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> Statistical and systematic errors combined.

<sup>2</sup> Statistical and systematic errors combined. Using values from fits with floating  $M2$  amplitudes  $a_2(\chi_{c1})$ ,  $a_2(\chi_{c2})$ ,  $b_2(\chi_{c1})$ ,  $b_2(\chi_{c2})$  and fixed  $E3$  amplitudes of  $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$ . Not independent of values for  $a_2(\chi_{c1}(1P))$  and  $a_2(\chi_{c2}(1P))$  from ARTUSO 09.

 **$b_2(\chi_{c2})/b_2(\chi_{c1})$  Magnetic quadrupole transition amplitude ratio**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>60 \pm 31</math> OUR AVERAGE</b>				
$74 \pm 40$	253k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$37^{+53}_{-47}$	59k	<sup>2</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> Statistical and systematic errors combined. Derived from the reported measurement of  $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$ .

<sup>2</sup> Statistical and systematic errors combined. Using values from fits with floating  $M2$  amplitudes  $a_2(\chi_{c1})$ ,  $a_2(\chi_{c2})$ ,  $b_2(\chi_{c1})$ ,  $b_2(\chi_{c2})$  and fixed  $E3$  amplitudes of  $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$ . Not independent of values for  $b_2(\chi_{c1}(1P))$  and  $b_2(\chi_{c2}(1P))$  from ARTUSO 09.

---

 **$\psi(2S)$  REFERENCES**

ABLIKIM	21AL PR D104 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AO PR D104 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21E PRL 126 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)



ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21S	PL B820 136576	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19AO	PR D99 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AT	PR D100 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18Q	PR D97 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18T	PR D98 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18Z	PL B783 452	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18	PL B781 174	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17X	PR D96 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOBBS	17	PR D96 092004	S. Dobbs <i>et al.</i>	(NWES, WAYN)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	16Y	JHEP 1605 132	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12Q	CP C36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)

ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
		Translated from ZETFP 85 429.		
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)

GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC
WIIK	75	Stanford Symp. 69	B.H. Wiik	(DESY)