

$\rho(2150)$ $I^G(J^{PC}) = 1^+(1^{--})$

OMITTED FROM SUMMARY TABLE

This entry was previously called $T_1(2190)$. See the review on "Spectroscopy of Light Meson Resonances."

 $\rho(2150)$ MASS**e⁺e⁻ PRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2034 ± 13 ± 9		1 ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\pi^0$
2111 ± 43 ± 25		2 ABLIKIM	21x BES3	$e^+e^- \rightarrow \eta'\pi^+\pi^-$
2255 ± 17 ± 50	1.8k	3 ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+K^-\eta$
2201 ± 19		4 LEES	20 BABR	$e^+e^- \rightarrow K^+K^-\gamma$
2227 ± 9 ± 9		5 LEES	20 RVUE	$e^+e^- \rightarrow K^+K^-$
2039 ± 8 ± 36		6 ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$
2239.2 $\pm 7.1 \pm 11.3$		7 ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2254 ± 22		8 LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
2150 ± 40 ± 50		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
1990 ± 80		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
2153 ± 37		BIAGINI	91 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, K^+K^-
2110 ± 50		9 CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-)$, $2(\pi^+\pi^-\pi^0)$

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\rho(770)$, $\rho(1450)$ and $\rho(1700)$. Could be another state.

² From a Breit-Wigner fit to the Born cross section, including an s -dependent continuum amplitude.

³ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (21.7 \pm 1.9^{+7.7}_{-8.3}) \times 10^{-6}$.

⁴ From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5σ .

⁵ From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

⁶ Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (6.7 \pm 1.1^{+2.2}_{-1.8}) \times 10^{-6}$.

⁷ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

⁸ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

⁹ Includes ATKINSON 85.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~2191	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~2070	¹ OAKDEN 94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2170	² MARTIN 80B	RVUE	
~2100	² MARTIN 80C	RVUE	

¹ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2110±35	¹ ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~2190	² CUTTS 78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155±15	^{2,3} COUPLAND 77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193± 2	^{2,4} ALSPECTOR 73	CNTR	$\bar{p}p$ S channel
2190±10	⁵ ABRAMS 70	CNTR	S channel $\bar{p}N$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

² Isospins 0 and 1 not separated.

³ From a fit to the total elastic cross section.

⁴ Referred to as T or T region by ALSPECTOR 73.

⁵ Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

 $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2140±30	ALDE 95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
2170±30	ALDE 92C	GAM4	100 $\pi^- p \rightarrow \omega\pi^0 n$

 $\rho(2150)$ WIDTH **$e^+ e^-$ PRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
234 ± 30 ± 25		¹ ABLIKIM 21A	BES3	$e^+ e^- \rightarrow \omega\pi^0$
135 ± 34 ± 30		² ABLIKIM 21X	BES3	$e^+ e^- \rightarrow \eta'\pi^+\pi^-$
460 ± 54 ± 160	1.8k	³ ABLIKIM 20F	BES3	$\psi(2S) \rightarrow K^+K^-\eta$
70 ± 38		⁴ LEES 20	BABR	$e^+ e^- \rightarrow K^+K^-\gamma$
127 ± 14 ± 4		⁵ LEES 20	RVUE	$e^+ e^- \rightarrow K^+K^-$
196 ± 23 ± 25		⁶ ABLIKIM 19AQ BES		$J/\psi \rightarrow K^+K^-\pi^0$
139.8 ± 12.3 ± 20.6		⁷ ABLIKIM 19L	BES3	$e^+ e^- \rightarrow K^+K^-$
109 ± 76		⁸ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+\pi^-\gamma$

350 \pm 40 \pm 50	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
310 \pm 140	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
389 \pm 79	BIAGINI	91 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $K^+ K^-$
410 \pm 100	⁹ CLEGG	90 RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-)$, $2(\pi^+ \pi^- \pi^0)$

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\rho(770)$, $\rho(1450)$ and $\rho(1700)$. Could be another state.

² From a Breit-Wigner fit to the Born cross section, including an s -dependent continuum amplitude.

³ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^-\eta = (21.7 \pm 1.9^{+7.7}_{-8.3}) \times 10^{-6}$.

⁴ From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5σ .

⁵ From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

⁶ Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+ K^-\pi^0 = (6.7 \pm 1.1^{+2.2}_{-1.8}) \times 10^{-6}$.

⁷ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

⁸ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

⁹ Includes ATKINSON 85.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
~ 296	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 40	¹ OAKDEN	94 RVUE	$0.36\text{--}1.55 \bar{p}p \rightarrow \pi\pi$
~ 250	² MARTIN	80B RVUE	
~ 200	² MARTIN	80C RVUE	

¹ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
230 ± 50	¹ ANISOVICH	02 SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135 ± 75	^{2,3} COUPLAND	77 CNTR	$0.7\text{--}2.4 \bar{p}p \rightarrow \bar{p}p$
98 ± 8	³ ALSPECTOR	73 CNTR	$\bar{p}p$ S channel
~ 85	⁴ ABRAMS	70 CNTR	S channel $\bar{p}N$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

² From a fit to the total elastic cross section.

³ Isospins 0 and 1 not separated.

⁴ Seen as bump in $J = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

$\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
320±70	ALDE	95 GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
~300	ALDE	92C GAM4	100 $\pi^- p \rightarrow \omega\pi^0 n$

 $\rho(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \pi^+ \pi^-$	seen
$\Gamma_3 K^+ K^-$	seen
$\Gamma_4 3(\pi^+ \pi^-)$	seen
$\Gamma_5 2(\pi^+ \pi^- \pi^0)$	seen
$\Gamma_6 \eta' \pi^+ \pi^-$	seen
$\Gamma_7 f_1(1285) \pi^+ \pi^-$	seen
$\Gamma_8 \omega \pi^0$	seen
$\Gamma_9 \omega \pi^0 \eta$	seen
$\Gamma_{10} p \bar{p}$	

 $\rho(2150) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\omega\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_8 \Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
34±11±16	ABLIKIM	21A BES3	$e^+ e^- \rightarrow \omega\pi^0$

$\Gamma(\eta' \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_6 \Gamma_1/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
23.3±5.3±3.3	¹ ABLIKIM	21X BES3	$e^+ e^- \rightarrow \eta' \pi^+ \pi^-$

¹ From a Breit-Wigner fit to the Born cross section interfering constructively with the continuum. For destructive interference the value is $0.64 \pm 0.49 \pm 0.42$ eV.

 $\rho(2150) \Gamma(i) \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(f_1(1285) \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$		
VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
3.1±0.6±0.5	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$

¹ Calculated by us from the reported value of cross section at the peak.

$\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma \times \Gamma_1/\Gamma$		
<u>VALUE (units 10^{-8})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.9 ± 1.9	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
¹ Calculated by us from the reported value of cross section at the peak.			

$\rho(2150)$ REFERENCES

ABLIKIM 21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 21X	PR D103 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES 20	PR D101 012011	J.P. Lees <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES 13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES 12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT 07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ALDE 95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ALDE 92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
BIAGINI 91	NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)
CLEGG 90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ATKINSON 85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE 75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER 68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)
GOUNARIS 68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	