



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

Neither of J or P has actually been measured.

Ξ_c^+ MASS

The fit uses the Ξ_c^+ and Ξ_c^0 mass and mass-difference measurements.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|----------|---|
| 2467.71 ± 0.23 OUR FIT | | Error includes scale factor of 1.3. | | |
| 2467.95 ± 0.19 OUR AVERAGE | | | | |
| 2467.97 ± 0.14 ± 0.17 | 3.8k | ¹ AAIJ | 14Z LHCb | pp at 7, 8 TeV |
| 2468.00 ± 0.18 ± 0.51 | 5.1k | AALTONEN | 14B CDF | $p\bar{p}$ at 1.96 TeV |
| 2468.1 ± 0.4 ⁺ 0.2 - 1.4 | 4.9k | ² LESIAK | 05 BELL | e^+e^- , $\Upsilon(4S)$ |
| 2465.8 ± 1.9 ± 2.5 | 90 | FRABETTI | 98 E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| 2467.0 ± 1.6 ± 2.0 | 147 | EDWARDS | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 2465.1 ± 3.6 ± 1.9 | 30 | ALBRECHT | 90F ARG | e^+e^- at $\Upsilon(4S)$ |
| 2467 ± 3 ± 4 | 23 | ALAM | 89 CLEO | e^+e^- 10.6 GeV |
| 2466.5 ± 2.7 ± 1.2 | 5 | BARLAG | 89C ACCM | π^- Cu 230 GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2464.4 ± 2.0 ± 1.4 | 30 | FRABETTI | 93B E687 | See FRABETTI 98 |
| 2459 ± 5 ± 30 | 56 | ³ COTEUS | 87 SPEC | $nA \simeq 600$ GeV |
| 2460 ± 25 | 82 | BIAGI | 83 SPEC | Σ^- Be 135 GeV |

¹ AAIJ 14Z systematic error includes in quadrature the 0.14 MeV uncertainty from the $m(\Lambda_c^+)$ mass value.

² The systematic error was (wrongly) given the other way round in LESIAK 05; see the erratum.

³ Although COTEUS 87 claims to agree well with BIAGI 83 on the mass and width, there appears to be a discrepancy between the two experiments. BIAGI 83 sees a single peak (stated significance about 6 standard deviations) in the $\Lambda K^- \pi^+ \pi^+$ mass spectrum. COTEUS 87 sees *two* peaks in the same spectrum, one at the Ξ_c^+ mass, the other 75 MeV lower. The latter is attributed to $\Xi_c^+ \rightarrow \Sigma^0 K^- \pi^+ \pi^+ \rightarrow (\Lambda \gamma) K^- \pi^+ \pi^+$, with the γ unseen. The *combined* significance of the double peak is stated to be 5.5 standard deviations. But the absence of any trace of a lower peak in BIAGI 83 seems to us to throw into question the interpretation of the lower peak of COTEUS 87.

Ξ_c^+ MEAN LIFE

| VALUE (10^{-15} s) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------------|-----------|--|
| 453 ± 5 OUR AVERAGE | | | | |
| 454 ± 5 ± 2 | 56k | ¹ AAIJ | 19AG LHCb | $\Xi_c^+ \rightarrow p K^- \pi^+$ |
| 503 ± 47 ± 18 | 250 | MAHMOOD | 02 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 439 ± 22 ± 9 | 532 | LINK | 01D FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

| | | | | | |
|-------------------------------|-----|----------|-----|------|---|
| $340_{-50}^{+70} \pm 20$ | 56 | FRABETTI | 98 | E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| $400_{-120}^{+180} \pm 100$ | 102 | COTEUS | 87 | SPEC | $nA \simeq 600$ GeV |
| $480_{-150}^{+210+200} - 100$ | 53 | BIAGI | 85C | SPEC | Σ^- Be 135 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------------------------|----|----------|-----|------|--------------------------|
| $410_{-80}^{+110} \pm 20$ | 30 | FRABETTI | 93B | E687 | See FRABETTI 98 |
| 200_{-60}^{+110} | 6 | BARLAG | 89C | ACCM | $\pi^- (K^-)$ Cu 230 GeV |

¹ AAIJ 19AG reports $[\Xi_C^+ \text{ MEAN LIFE}] / [D^\pm \text{ MEAN LIFE}] = 0.4392 \pm 0.0034 \pm 0.0028$
 which we multiply by our best value $D^\pm \text{ MEAN LIFE} = (1.033 \pm 0.005) \times 10^{-12}$ s.
 Our first error is their experiment's error and our second error is the systematic error from using our best value.

Ξ_C^+ DECAY MODES

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Xi_C^+ \rightarrow \Sigma^+ \bar{K}^*(892)^0$ seen in $\Xi_C^+ \rightarrow \Sigma^+ K^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|------|--------------------------------|-----------------------------------|
|------|--------------------------------|-----------------------------------|

Cabibbo-favored ($S = -2$) decays

| | | | |
|---------------|----------------------------------|--------------------------------|--------|
| Γ_1 | $p2K_S^0$ | $(2.5 \pm 1.3) \times 10^{-3}$ | |
| Γ_2 | $\Lambda \bar{K}^0 \pi^+$ | — | |
| Γ_3 | $\Sigma(1385)^+ \bar{K}^0$ | [a] $(2.9 \pm 2.0) \%$ | |
| Γ_4 | $\Lambda K^- 2\pi^+$ | $(9 \pm 4) \times 10^{-3}$ | |
| Γ_5 | $\Lambda \bar{K}^*(892)^0 \pi^+$ | [a] $< 5 \times 10^{-3}$ | CL=90% |
| Γ_6 | $\Sigma(1385)^+ K^- \pi^+$ | [a] $< 6 \times 10^{-3}$ | CL=90% |
| Γ_7 | $\Sigma^+ K^- \pi^+$ | $(2.7 \pm 1.2) \%$ | |
| Γ_8 | $\Sigma^+ \bar{K}^*(892)^0$ | [a] $(2.3 \pm 1.1) \%$ | |
| Γ_9 | $\Sigma^0 K^- 2\pi^+$ | $(8 \pm 5) \times 10^{-3}$ | |
| Γ_{10} | $\Xi^0 \pi^+$ | $(1.6 \pm 0.8) \%$ | |
| Γ_{11} | $\Xi^- 2\pi^+$ | $(2.9 \pm 1.3) \%$ | |
| Γ_{12} | $\Xi(1530)^0 \pi^+$ | [a] $< 2.9 \times 10^{-3}$ | CL=90% |
| Γ_{13} | $\Xi(1620)^0 \pi^+$ | seen | |
| Γ_{14} | $\Xi(1690)^0 \pi^+$ | seen | |
| Γ_{15} | $\Xi^0 \pi^+ \pi^0$ | $(6.7 \pm 3.5) \%$ | |
| Γ_{16} | $\Xi^0 \pi^- 2\pi^+$ | $(5.0 \pm 2.6) \%$ | |
| Γ_{17} | $\Xi^0 e^+ \nu_e$ | $(7 \pm 4) \%$ | |
| Γ_{18} | $\Omega^- K^+ \pi^+$ | $(2.0 \pm 1.5) \times 10^{-3}$ | |

Cabibbo-suppressed decays

| | | | |
|---------------|---|------------------------------------|---------|
| Γ_{19} | $\rho K^- \pi^+$ | $(6.2 \pm 3.0) \times 10^{-3}$ | $S=1.5$ |
| Γ_{20} | $\rho \bar{K}^*(892)^0$ | [a] $(3.3 \pm 1.7) \times 10^{-3}$ | |
| Γ_{21} | $\Sigma^+ \pi^+ \pi^-$ | $(1.4 \pm 0.8) \%$ | |
| Γ_{22} | $\Sigma^- 2\pi^+$ | $(5.1 \pm 3.4) \times 10^{-3}$ | |
| Γ_{23} | $\Sigma^+ K^+ K^-$ | $(4.3 \pm 2.5) \times 10^{-3}$ | |
| Γ_{24} | $\Sigma^+ \phi$ | [a] $< 3.2 \times 10^{-3}$ | CL=90% |
| Γ_{25} | $\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$ | $< 1.3 \times 10^{-3}$ | CL=90% |
| Γ_{26} | $\rho \phi(1020)$ | $(1.2 \pm 0.6) \times 10^{-4}$ | |

[a] This branching fraction includes all the decay modes of the final-state resonance.

 Ξ_c^+ BRANCHING RATIOS**Cabibbo-favored ($S = -2$) decays**

$\Gamma(\rho 2K_S^0)/\Gamma(\Xi^- 2\pi^+)$ Γ_1/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|-------------|------|-------------------------------|
| $0.087 \pm 0.016 \pm 0.014$ | 168 ± 27 | LESLIAK | 05 | BELL $e^+ e^-$, $\gamma(4S)$ |

$\Gamma(\Sigma(1385)^+ \bar{K}^0)/\Gamma(\Xi^- 2\pi^+)$ Γ_3/Γ_{11}

Unseen decay modes of the $\Sigma(1385)^+$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|------------------------|
| $1.00 \pm 0.49 \pm 0.24$ | 20 | LINK | 03E | FOCS < 1.72 , 90% CL |

$\Gamma(\Lambda K^- 2\pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_4/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|---------------|-------------|------|---|
| 0.323 ± 0.033 OUR AVERAGE | | | | |
| $0.32 \pm 0.03 \pm 0.02$ | 1177 ± 55 | LESLIAK | 05 | BELL $e^+ e^-$, $\gamma(4S)$ |
| $0.28 \pm 0.06 \pm 0.06$ | 58 | LINK | 03E | FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| $0.58 \pm 0.16 \pm 0.07$ | 61 | BERGFELD | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Lambda \bar{K}^*(892)^0 \pi^+)/\Gamma(\Lambda K^- 2\pi^+)$ Γ_5/Γ_4

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|-----------------------------------|
| < 0.5 | 90 | BERGFELD | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma(1385)^+ K^- \pi^+)/\Gamma(\Lambda K^- 2\pi^+)$ Γ_6/Γ_4

Unseen decay modes of the $\Sigma(1385)^+$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|-----------------------------------|
| < 0.7 | 90 | BERGFELD | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^+ K^- \pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_7/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|------------------|------|---|
| 0.94 ± 0.10 OUR AVERAGE | | | | |
| $0.91 \pm 0.11 \pm 0.04$ | 251 | LINK | 03E | FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| $0.92 \pm 0.20 \pm 0.07$ | | ¹ JUN | 00 | SELX Σ^- nucleus, 600 GeV |
| $1.18 \pm 0.26 \pm 0.17$ | 119 | BERGFELD | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |

¹ This JUN 00 result is redundant with other results given below.

$\Gamma(\Sigma^+ \bar{K}^*(892)^0)/\Gamma(\Xi^- 2\pi^+)$ Γ_8/Γ_{11} Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--|
| 0.81±0.15 OUR AVERAGE | | | | |
| 0.78±0.16±0.06 | 119 | LINK | 03E FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.92±0.27±0.14 | 61 | BERGFELD | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

 $\Gamma(\Sigma^0 K^- 2\pi^+)/\Gamma(\Lambda K^- 2\pi^+)$ Γ_9/Γ_4

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------|------|---------------------|---------|----------------------|
| 0.84±0.36 | 47 | ¹ COTEUS | 87 SPEC | $nA \approx 600$ GeV |

¹ See, however, the note on the COTEUS 87 Ξ_c^+ mass measurement. $\Gamma(\Xi^0 \pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{10}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|-------------------------------|
| 0.55±0.13±0.09 | 39 | EDWARDS | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

 $\Gamma(\Xi^- 2\pi^+)/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-----------------|----------|-------------------------------|
| 2.86±1.21±0.38 | 24 | ¹ LI | 19C BELL | $e^+e^- \approx \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|-----|----------|----------|---|
| seen | 131 | BERGFELD | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| seen | 160 | AVERY | 95 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| seen | 30 | FRABETTI | 93B E687 | γ Be, $\bar{E}_\gamma = 220$ GeV |
| seen | 30 | ALBRECHT | 90F ARG | e^+e^- at $\Upsilon(4S)$ |
| seen | 23 | ALAM | 89 CLEO | e^+e^- 10.6 GeV |

¹ LI 19C report a significance of 6.8 σ for the observation of this decay mode, observed in Ξ_c^+ from $B^0 \rightarrow \bar{\Lambda}_c^- \Xi_c^+$. $\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{12}/Γ_{11} Unseen decay modes of the $\Xi(1530)^0$ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|-------------|----------|--|
| <0.1 | 90 | LINK | 03E FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|----------|---------|-------------------------------|
| <0.2 | 90 | BERGFELD | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
|------|----|----------|---------|-------------------------------|

 $\Gamma(\Xi(1620)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------------|
| seen | SUMIHAMA 19 | BELL | e^+e^- mostly at $\Upsilon(4S)$ |

 $\Gamma(\Xi(1690)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{14}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------------|
| seen | SUMIHAMA 19 | BELL | e^+e^- mostly at $\Upsilon(4S)$ |

 $\Gamma(\Xi^0 \pi^+ \pi^0)/\Gamma(\Xi^- 2\pi^+)$ Γ_{15}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|-------------------------------|
| 2.34±0.57±0.37 | 81 | EDWARDS | 96 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^0 \pi^+ \pi^0)$ Γ_{12}/Γ_{15}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|---------|----|-----------------------------------|
| <0.3 | 90 | EDWARDS | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |
|------|----|---------|----|-----------------------------------|

 $\Gamma(\Xi^0 \pi^- 2\pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{16}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|---------------------------|----|---------|----|-----------------------------------|
| 1.74 ± 0.42 ± 0.27 | 57 | EDWARDS | 96 | CLE2 $e^+ e^- \approx \gamma(4S)$ |
|---------------------------|----|---------|----|-----------------------------------|

 $\Gamma(\Xi^0 e^+ \nu_e)/\Gamma(\Xi^- 2\pi^+)$ Γ_{17}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|--|----|-----------|-----|-----------------------------------|
| 2.3 ± 0.6^{+0.3}_{-0.6} | 41 | ALEXANDER | 95B | CLE2 $e^+ e^- \approx \gamma(4S)$ |
|--|----|-----------|-----|-----------------------------------|

 $\Gamma(\Omega^- K^+ \pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{18}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|---------------------------|----|------|-----|---------------------|
| 0.07 ± 0.03 ± 0.03 | 14 | LINK | 03E | FOCS < 0.12, 90% CL |
|---------------------------|----|------|-----|---------------------|

 $\Gamma(p K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|------------------------------|-------|-------------------------------|--|--|
| 6.2 ± 3.0 OUR AVERAGE | Error | includes scale factor of 1.5. | | |
|------------------------------|-------|-------------------------------|--|--|

| | | | | |
|---------------------|------|-------------------|------|---------------------|
| 11.35 ± 0.02 ± 3.87 | 1.6M | ¹ AAIJ | 20AH | LHCB pp at 13 TeV |
|---------------------|------|-------------------|------|---------------------|

| | | | | |
|-----------------|----|-----------------|-----|-----------------------------------|
| 4.5 ± 2.1 ± 0.7 | 24 | ² LI | 19C | BELL $e^+ e^- \approx \gamma(4S)$ |
|-----------------|----|-----------------|-----|-----------------------------------|

¹AAIJ 20AH extracts $B(\Xi_c^+ \rightarrow p K^- \pi^+)$ assuming production fraction ratios $f_{\Xi_c^0}/f_{\Lambda_c^+} = (9.7 \pm 0.9 \pm 3.1) \times 10^{-2}$ (from AAIJ 19AB plus heavy quark symmetry arguments) as well as $f_{\Xi_c^0}/f_{\Xi_c^+} = 1.00 \pm 0.01$, and uses the input $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.23 \pm 0.33) \times 10^{-2}$. Its correlation with $B(\Xi_c^0 \rightarrow \Lambda_c^+ \pi^-)$, as measured in AAIJ 20AH, is 0.414.

²LI 19C report a significance of 4.4σ for the observation of this decay mode, observed in Ξ_c^+ from $\bar{B}^0 \rightarrow \bar{\Lambda}_c^+ \Xi_c^+$.

———— Cabibbo-suppressed decays ————

 $\Gamma(p K^- \pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{19}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|--------------------------------|--|--|--|--|
| 0.21 ± 0.04 OUR AVERAGE | | | | |
|--------------------------------|--|--|--|--|

| | | | | |
|---------------|---------|----------------|------|-----------------------------|
| 0.194 ± 0.054 | 47 ± 11 | VAZQUEZ-JA..08 | SELX | Σ^- nucleus, 600 GeV |
|---------------|---------|----------------|------|-----------------------------|

| | | | | |
|-----------------------|-----|------|-----|-----------------------|
| 0.234 ± 0.047 ± 0.022 | 202 | LINK | 01B | FOCS γ nucleus |
|-----------------------|-----|------|-----|-----------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|----|-----|----|------------------------------|
| 0.20 ± 0.04 ± 0.02 | 76 | JUN | 00 | SELX See VAZQUEZ-JAUREGUI 08 |
|--------------------|----|-----|----|------------------------------|

 $\Gamma(p \bar{K}^*(892)^0)/\Gamma(p K^- \pi^+)$ Γ_{20}/Γ_{19} Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|---------------------------|------|-----|-----------------------|
| 0.54 ± 0.09 ± 0.05 | LINK | 01B | FOCS γ nucleus |
|---------------------------|------|-----|-----------------------|

$\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma(\Xi^- 2\pi^+)$ Γ_{21}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------|--------|-----------------|------|-----------------------------|
| 0.48±0.20 | 21 ± 8 | VAZQUEZ-JA...08 | SELX | Σ^- nucleus, 600 GeV |

 $\Gamma(\Sigma^- 2\pi^+)/\Gamma(\Xi^- 2\pi^+)$ Γ_{22}/Γ_{11}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------|--------|-----------------|------|-----------------------------|
| 0.18±0.09 | 10 ± 4 | VAZQUEZ-JA...08 | SELX | Σ^- nucleus, 600 GeV |

 $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ K^- \pi^+)$ Γ_{23}/Γ_7

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.16±0.06±0.01 | 17 | LINK | 03E FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

 $\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ K^- \pi^+)$ Γ_{24}/Γ_7 Unseen decay modes of the ϕ are included.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|----------|--|
| <0.12 | 90 | LINK | 03E FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

 $\Gamma(p\phi(1020))/\Gamma(pK^- \pi^+)$ Γ_{26}/Γ_{19}

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------------|----------|---------------|
| 19.8±0.7±0.9±0.2 | 3.4k | ¹ AAIJ | 19i LHCb | pp at 8 TeV |

¹ The last uncertainty is due to the uncertainty in the $\phi \rightarrow K^+ K^-$ branching fraction. $\Gamma(\Xi(1690)^0 K^+ \times B(\Xi(1690)^0 \rightarrow \Sigma^+ K^-))/\Gamma(\Sigma^+ K^- \pi^+)$ Γ_{25}/Γ_7

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|----------|--|
| <0.05 | 90 | LINK | 03E FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

 Ξ_c^+ REFERENCES

| | | | | |
|---------------|------|-----------------------|-----------------------------------|----------------------|
| AAIJ | 20AH | PR D102 071101 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 19AB | PR D99 052006 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 19AG | PR D100 032001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 19I | JHEP 1904 084 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| LI | 19C | PR D100 031101 | Y.B. Li <i>et al.</i> | (BELLE Collab.) |
| SUMIHAMA | 19 | PRL 122 072501 | M. Sumihama <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 14Z | PRL 113 032001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 14B | PR D89 072014 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| VAZQUEZ-JA... | 08 | PL B666 299 | E. Vazquez-Jauregui <i>et al.</i> | (SELEX Collab.) |
| LESIK | 05 | PL B605 237 | T. Lesiak <i>et al.</i> | (BELLE Collab.) |
| Also | | PL B617 198 (erratum) | T. Lesiak <i>et al.</i> | (BELLE Collab.) |
| LINK | 03E | PL B571 139 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| MAHMOOD | 02 | PR D65 031102 | A.H. Mahmood <i>et al.</i> | (CLEO Collab.) |
| LINK | 01B | PL B512 277 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| LINK | 01D | PL B523 53 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| JUN | 00 | PRL 84 1857 | S.Y. Jun <i>et al.</i> | (FNAL SELEX Collab.) |
| FRABETTI | 98 | PL B427 211 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| BERGFELD | 96 | PL B365 431 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 96 | PL B373 261 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| ALEXANDER | 95B | PRL 74 3113 | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| Also | | PRL 75 4155 (erratum) | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| AVERY | 95 | PRL 75 4364 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 93B | PRL 70 1381 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| ALBRECHT | 90F | PL B247 121 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALAM | 89 | PL B226 401 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| BARLAG | 89C | PL B233 522 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| COTEUS | 87 | PRL 59 1530 | P. Coteus <i>et al.</i> | (FNAL E400 Collab.) |
| BIAGI | 85C | PL 150B 230 | S.F. Biagi <i>et al.</i> | (CERN WA62 Collab.) |
| BIAGI | 83 | PL 122B 455 | S.F. Biagi <i>et al.</i> | (CERN WA62 Collab.) |